

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex**Session Type:** Rate**Suggested Citation****Software:** Surveillance Research Program, National Cancer Institute SEER*Stat software (www.seer.cancer.gov/seerstat) version 6.2.3.**Data:** Alaska Cancer Registry SEER*Stat Database - Incidence data for diagnosis years 1996 to 2003. Released December 2006. Alaska population data for rate calculations are from the SEER program (<http://www.seer.cancer.gov/popdata/>) and include populations by borough/census area, expanded races, and Hispanic origin.**Data****Database:** ACR w/Borough, Race, & Hispanic Pops (1996-2003), December 2005**Statistic****Statistic:** Age-Adjusted Rates (Age recode with <1 year olds) (2000 US Std Population (19 age groups - Census P25-1130) standard)**P-Values:** .05**Options:** Display Confidence Intervals and Standard Errors**Selection****Option:** Select only malignant behavior**Table****Page:** Year of diagnosis**Row:** Site rec with Kaposi and mesothelioma**Column:** Race recode Y - 4 groups only [Race recode Y]
Sex - M & F only [Sex]**Output****Title:** Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex**Suppress:** Statistics based on fewer than 6 cases**Display Rates as:** Cases per 100,000**Precision:** 1 decimal place (default)

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

| | Total | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | White | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 562.4 | 546.4 | 578.8 | 6,550 | 429.4 | 417.4 | 441.8 | 5,592 |
| Oral Cavity and Pharynx | 14.4 | 12.2 | 17.3 | 208 | 6.7 | 5.2 | 8.5 | 77 |
| Lip | 1.8 | 0.9 | 3.6 | 17 | 0.5 | 0.2 | 1.3 | 6 |
| Tongue | 4.1 | 2.9 | 5.9 | 58 | 1.6 | 0.9 | 2.6 | 18 |
| Salivary Gland | 1.5 | 0.8 | 2.9 | 17 | 0.9 | 0.4 | 1.7 | 10 |
| Floor of Mouth | 1.2 | 0.7 | 2.4 | 19 | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | 0.9 | 0.5 | 2.0 | 15 | 1.4 | 0.8 | 2.5 | 15 |
| Nasopharynx | 0.7 | 0.3 | 1.9 | 11 | 0.6 | 0.3 | 1.4 | 8 |
| Tonsil | 2.4 | 1.6 | 3.8 | 42 | ^ | ^ | ^ | ^ |
| Oropharynx | 0.7 | 0.2 | 1.9 | 9 | ^ | ^ | ^ | ^ |
| Hypopharynx | 1.0 | 0.5 | 2.2 | 14 | ^ | ^ | ^ | ^ |
| Other Oral Cavity and Pharynx | 0.3 | 0.1 | 1.4 | 6 | ^ | ^ | ^ | ^ |
| Digestive System | 98.4 | 91.7 | 105.8 | 1,115 | 68.0 | 62.9 | 73.4 | 751 |
| Esophagus | 8.4 | 6.6 | 10.8 | 103 | 2.3 | 1.4 | 3.4 | 26 |
| Stomach | 8.3 | 6.3 | 11.0 | 86 | 3.6 | 2.5 | 5.1 | 40 |
| Small Intestine | 1.5 | 0.8 | 2.9 | 19 | 1.3 | 0.7 | 2.4 | 14 |
| Colon and Rectum | 57.2 | 52.0 | 62.9 | 620 | 40.7 | 36.7 | 45.0 | 440 |
| Colon excluding Rectum | 41.7 | 37.1 | 46.9 | 413 | 29.1 | 25.7 | 32.8 | 303 |
| Cecum | 7.7 | 5.7 | 10.4 | 72 | 9.1 | 7.3 | 11.4 | 90 |
| Appendix | 0.5 | 0.3 | 1.6 | 11 | ^ | ^ | ^ | ^ |
| Ascending Colon | 9.1 | 6.9 | 11.9 | 78 | 4.8 | 3.5 | 6.5 | 46 |
| Hepatic Flexure | 1.7 | 1.0 | 3.2 | 20 | 1.9 | 1.1 | 3.0 | 19 |
| Transverse Colon | 2.8 | 1.9 | 4.5 | 33 | 1.8 | 1.0 | 2.9 | 19 |
| Splenic Flexure | 1.0 | 0.5 | 2.2 | 12 | 0.9 | 0.4 | 1.8 | 10 |
| Descending Colon | 3.6 | 2.4 | 5.6 | 34 | 1.2 | 0.6 | 2.2 | 13 |
| Sigmoid Colon | 11.9 | 9.6 | 14.9 | 121 | 7.4 | 5.8 | 9.3 | 83 |
| Large Intestine, NOS | 3.3 | 2.1 | 5.2 | 32 | 1.8 | 1.1 | 2.9 | 21 |
| Rectum and Rectosigmoid Junction | 15.5 | 13.2 | 18.4 | 207 | 11.6 | 9.6 | 13.9 | 137 |
| Rectosigmoid Junction | 4.4 | 3.2 | 6.2 | 56 | 2.1 | 1.3 | 3.2 | 26 |
| Rectum | 11.1 | 9.2 | 13.6 | 151 | 9.5 | 7.7 | 11.6 | 111 |
| Anus, Anal Canal and Anorectum | 0.5 | 0.2 | 1.5 | 10 | 2.5 | 1.7 | 3.6 | 34 |
| Liver and Intrahepatic Bile Duct | 8.1 | 6.4 | 10.5 | 109 | 3.3 | 2.3 | 4.6 | 42 |
| Liver | 7.2 | 5.6 | 9.5 | 98 | 2.6 | 1.7 | 3.8 | 32 |
| Intrahepatic Bile Duct | 0.9 | 0.3 | 2.3 | 11 | 0.7 | 0.3 | 1.5 | 10 |
| Gallbladder | 0.9 | 0.4 | 2.2 | 10 | 0.8 | 0.3 | 1.7 | 7 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

| | Total | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Black | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 651.8 | 558.2 | 770.1 | 241 | 407.5 | 340.7 | 486.5 | 180 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Tongue | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 50.6 | 0 | ^ | ^ | ^ | ^ |
| Floor of Mouth | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 20.1 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 20.1 | 0 |
| Tonsil | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Oropharynx | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Hypopharynx | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Digestive System | 152.8 | 104.8 | 230.5 | 51 | 76.7 | 47.4 | 120.1 | 26 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | 28.3 | 10.2 | 84.2 | 7 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 20.1 | 0 |
| Colon and Rectum | 71.9 | 38.7 | 138.7 | 26 | 67.9 | 39.9 | 110.5 | 21 |
| Colon excluding Rectum | 51.3 | 28.4 | 107.9 | 22 | 48.4 | 25.5 | 86.1 | 15 |
| Cecum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 50.6 | 0 | ^ | ^ | ^ | ^ |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 20.1 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Descending Colon | 0.0 | 0.0 | 50.6 | 0 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | 19.5 | 6.2 | 49.1 | 6 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 20.1 | 0 |
| Rectum | ^ | ^ | ^ | ^ | 19.5 | 6.2 | 49.1 | 6 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 20.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

| | Total | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | American Indian/Alaska Native | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 557.0 | 520.9 | 595.6 | 1,137 | 484.1 | 456.4 | 513.3 | 1,257 |
| Oral Cavity and Pharynx | 26.1 | 19.6 | 35.3 | 62 | 10.3 | 6.6 | 15.6 | 26 |
| Lip | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Tongue | 3.6 | 1.5 | 9.3 | 8 | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | 1.8 | 0.7 | 4.8 | 6 |
| Floor of Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | 2.5 | 0.8 | 7.8 | 6 | ^ | ^ | ^ | ^ |
| Nasopharynx | 11.0 | 6.9 | 18.2 | 25 | 4.2 | 1.8 | 8.4 | 9 |
| Tonsil | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Oropharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Digestive System | 204.1 | 181.8 | 229.1 | 391 | 162.7 | 146.2 | 180.8 | 385 |
| Esophagus | 13.7 | 8.5 | 22.0 | 25 | 6.2 | 3.3 | 10.8 | 14 |
| Stomach | 31.4 | 23.4 | 42.2 | 65 | 18.5 | 13.4 | 25.3 | 47 |
| Small Intestine | ^ | ^ | ^ | ^ | 2.6 | 1.0 | 6.0 | 7 |
| Colon and Rectum | 111.7 | 95.5 | 130.7 | 214 | 101.1 | 88.1 | 115.7 | 235 |
| Colon excluding Rectum | 78.9 | 64.9 | 95.7 | 142 | 75.4 | 64.2 | 88.3 | 172 |
| Cecum | 12.8 | 7.7 | 21.0 | 24 | 16.0 | 11.0 | 22.7 | 35 |
| Appendix | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Ascending Colon | 8.5 | 4.4 | 16.0 | 15 | 8.4 | 4.9 | 13.6 | 19 |
| Hepatic Flexure | 4.2 | 1.5 | 10.5 | 8 | 6.0 | 3.0 | 10.8 | 12 |
| Transverse Colon | 10.4 | 5.6 | 18.7 | 17 | 12.7 | 8.2 | 19.1 | 26 |
| Splenic Flexure | ^ | ^ | ^ | ^ | 3.1 | 1.3 | 6.8 | 8 |
| Descending Colon | ^ | ^ | ^ | ^ | 2.5 | 0.8 | 6.1 | 6 |
| Sigmoid Colon | 32.9 | 24.5 | 44.2 | 63 | 24.1 | 18.2 | 31.6 | 60 |
| Large Intestine, NOS | 4.8 | 1.5 | 12.1 | 6 | 2.6 | 0.9 | 6.3 | 6 |
| Rectum and Rectosigmoid Junction | 32.9 | 25.0 | 43.5 | 72 | 25.7 | 19.5 | 33.5 | 63 |
| Rectosigmoid Junction | 8.1 | 4.6 | 14.7 | 18 | 5.9 | 3.1 | 10.5 | 14 |
| Rectum | 24.8 | 17.9 | 34.5 | 54 | 19.8 | 14.4 | 26.8 | 49 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | 3.2 | 1.3 | 7.0 | 8 |
| Liver and Intrahepatic Bile Duct | 12.3 | 7.2 | 20.6 | 24 | 5.5 | 2.8 | 9.9 | 14 |
| Liver | 9.6 | 5.2 | 17.3 | 20 | 5.0 | 2.5 | 9.3 | 13 |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gallbladder | 6.8 | 2.8 | 14.5 | 9 | 5.3 | 2.6 | 9.9 | 11 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

| | Total | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Asian or Pacific Islander | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 437.8 | 377.8 | 507.7 | 244 | 285.0 | 248.4 | 328.7 | 279 |
| Oral Cavity and Pharynx | 9.5 | 3.4 | 29.0 | 7 | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Tongue | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 13.8 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 13.8 | 0 |
| Oropharynx | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Hypopharynx | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 13.8 | 0 |
| Digestive System | 133.5 | 102.4 | 174.8 | 79 | 74.1 | 54.6 | 101.6 | 63 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | 31.7 | 16.8 | 58.7 | 16 | 11.9 | 5.4 | 28.0 | 10 |
| Small Intestine | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Colon and Rectum | 31.5 | 18.0 | 56.5 | 19 | 30.6 | 19.2 | 50.6 | 28 |
| Colon excluding Rectum | 21.0 | 10.3 | 43.9 | 13 | 17.4 | 10.0 | 33.8 | 17 |
| Cecum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 17.8 | 0 | ^ | ^ | ^ | ^ |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Descending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 13.8 | 0 |
| Sigmoid Colon | ^ | ^ | ^ | ^ | 8.0 | 3.4 | 22.7 | 8 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 10.4 | 3.6 | 30.6 | 6 | 13.2 | 5.5 | 30.9 | 11 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | ^ | ^ | ^ | ^ | 11.2 | 4.0 | 28.7 | 9 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Liver and Intrahepatic Bile Duct | 39.6 | 24.8 | 65.4 | 27 | 12.7 | 6.0 | 28.9 | 11 |
| Liver | 38.9 | 24.2 | 64.6 | 26 | 12.7 | 6.0 | 28.9 | 11 |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 13.8 | 0 |
| Gallbladder | 0.0 | 0.0 | 17.8 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

| | Total | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | White | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 1.2 | 0.6 | 2.6 | 14 | 1.4 | 0.7 | 2.5 | 14 |
| Pancreas | 11.4 | 9.2 | 14.3 | 131 | 10.3 | 8.4 | 12.6 | 114 |
| Retroperitoneum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Peritoneum, Omentum and Mesentery | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Digestive Organs | 0.7 | 0.3 | 2.0 | 8 | 1.3 | 0.6 | 2.3 | 12 |
| Respiratory System | 94.1 | 87.6 | 101.2 | 1,022 | 64.9 | 60.0 | 70.1 | 734 |
| Nose, Nasal Cavity and Middle Ear | 0.8 | 0.3 | 2.0 | 10 | ^ | ^ | ^ | ^ |
| Larynx | 6.6 | 5.0 | 8.7 | 82 | 1.4 | 0.9 | 2.4 | 20 |
| Lung and Bronchus | 86.6 | 80.3 | 93.5 | 926 | 62.9 | 58.2 | 68.1 | 706 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Bones and Joints | 0.9 | 0.6 | 2.0 | 20 | 0.9 | 0.5 | 1.6 | 16 |
| Soft Tissue including Heart | 3.5 | 2.4 | 5.2 | 51 | 2.6 | 1.8 | 3.7 | 40 |
| Skin excluding Basal and Squamous | 17.6 | 15.2 | 20.7 | 267 | 12.7 | 10.9 | 14.8 | 200 |
| Melanoma of the Skin | 16.4 | 14.0 | 19.3 | 249 | 11.6 | 9.8 | 13.6 | 185 |
| Other Non-Epithelial Skin | 1.3 | 0.7 | 2.6 | 18 | 1.1 | 0.6 | 2.0 | 15 |
| Breast | 0.9 | 0.4 | 2.2 | 12 | 141.0 | 134.4 | 147.9 | 1,992 |
| Female Genital System | 0.0 | 0.0 | 1.1 | 0 | 48.6 | 44.8 | 52.8 | 682 |
| Cervix Uteri | 0.0 | 0.0 | 1.1 | 0 | 8.4 | 7.0 | 10.1 | 141 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 1.1 | 0 | 22.4 | 19.8 | 25.3 | 306 |
| Corpus Uteri | 0.0 | 0.0 | 1.1 | 0 | 21.3 | 18.7 | 24.1 | 292 |
| Uterus, NOS | 0.0 | 0.0 | 1.1 | 0 | 1.1 | 0.6 | 2.1 | 14 |
| Ovary | 0.0 | 0.0 | 1.1 | 0 | 13.6 | 11.5 | 16.0 | 181 |
| Vagina | 0.0 | 0.0 | 1.1 | 0 | 0.8 | 0.3 | 1.6 | 9 |
| Vulva | 0.0 | 0.0 | 1.1 | 0 | 2.7 | 1.8 | 3.9 | 34 |
| Other Female Genital Organs | 0.0 | 0.0 | 1.1 | 0 | 0.8 | 0.4 | 1.6 | 11 |
| Male Genital System | 184.1 | 175.1 | 193.7 | 2,070 | 0.0 | 0.0 | 0.5 | 0 |
| Prostate | 176.6 | 167.7 | 186.1 | 1,923 | 0.0 | 0.0 | 0.5 | 0 |
| Testis | 6.4 | 5.3 | 8.0 | 134 | 0.0 | 0.0 | 0.5 | 0 |
| Penis | 0.9 | 0.3 | 2.3 | 10 | 0.0 | 0.0 | 0.5 | 0 |
| Other Male Genital Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 0.5 | 0 |
| Urinary System | 65.3 | 59.7 | 71.5 | 698 | 18.4 | 15.9 | 21.3 | 217 |
| Urinary Bladder | 46.7 | 41.8 | 52.3 | 454 | 9.8 | 7.9 | 12.0 | 106 |
| Kidney and Renal Pelvis | 17.3 | 14.8 | 20.4 | 232 | 8.1 | 6.5 | 10.0 | 105 |
| Ureter | 1.0 | 0.4 | 2.3 | 9 | ^ | ^ | ^ | ^ |
| Other Urinary Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

| | Total | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Black | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Pancreas | 18.7 | 5.7 | 71.2 | 6 | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Peritoneum, Omentum and Mesentery | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 20.1 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 20.1 | 0 |
| Respiratory System | 116.5 | 78.6 | 184.4 | 39 | 53.0 | 31.4 | 88.1 | 21 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 50.6 | 0 | ^ | ^ | ^ | ^ |
| Larynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 104.5 | 69.2 | 170.1 | 36 | 48.5 | 27.8 | 82.8 | 19 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 20.1 | 0 |
| Bones and Joints | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Breast | ^ | ^ | ^ | ^ | 127.3 | 92.9 | 174.1 | 64 |
| Female Genital System | 0.0 | 0.0 | 50.6 | 0 | 46.8 | 27.8 | 78.8 | 24 |
| Cervix Uteri | 0.0 | 0.0 | 50.6 | 0 | 12.1 | 4.9 | 33.9 | 8 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 50.6 | 0 | 18.9 | 6.5 | 46.9 | 8 |
| Corpus Uteri | 0.0 | 0.0 | 50.6 | 0 | 18.9 | 6.5 | 46.9 | 8 |
| Uterus, NOS | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Ovary | 0.0 | 0.0 | 50.6 | 0 | 11.0 | 3.8 | 33.3 | 6 |
| Vagina | 0.0 | 0.0 | 50.6 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Male Genital System | 267.6 | 212.1 | 349.5 | 100 | 0.0 | 0.0 | 20.1 | 0 |
| Prostate | 266.8 | 211.4 | 348.7 | 99 | 0.0 | 0.0 | 20.1 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 20.1 | 0 |
| Penis | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Urinary System | 39.9 | 19.3 | 95.6 | 14 | 39.4 | 18.5 | 75.9 | 11 |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 26.6 | 10.7 | 80.0 | 10 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 50.6 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

| | Total | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | American Indian/Alaska Native | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 2.8 | 0.9 | 8.5 | 6 | 4.2 | 2.0 | 8.3 | 10 |
| Pancreas | 20.1 | 13.4 | 29.9 | 37 | 14.3 | 9.8 | 20.4 | 35 |
| Retroperitoneum | 0.0 | 0.0 | 4.9 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Peritoneum, Omentum and Mesentery | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 130.7 | 113.9 | 150.1 | 263 | 72.7 | 61.7 | 85.2 | 167 |
| Nose, Nasal Cavity and Middle Ear | 3.0 | 0.9 | 8.8 | 6 | ^ | ^ | ^ | ^ |
| Larynx | 4.7 | 2.4 | 10.3 | 12 | 2.3 | 0.9 | 5.7 | 6 |
| Lung and Bronchus | 122.3 | 106.0 | 141.2 | 243 | 68.6 | 57.9 | 80.8 | 157 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | 1.4 | 0.5 | 6.2 | 6 | 2.4 | 0.9 | 5.8 | 8 |
| Skin excluding Basal and Squamous | 3.1 | 1.0 | 8.8 | 6 | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | 0.0 | 0.0 | 4.9 | 0 | 135.6 | 121.9 | 150.7 | 385 |
| Female Genital System | 0.0 | 0.0 | 4.9 | 0 | 31.4 | 25.0 | 39.4 | 90 |
| Cervix Uteri | 0.0 | 0.0 | 4.9 | 0 | 8.5 | 5.4 | 13.1 | 27 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 4.9 | 0 | 12.7 | 8.7 | 18.3 | 35 |
| Corpus Uteri | 0.0 | 0.0 | 4.9 | 0 | 12.4 | 8.4 | 17.9 | 34 |
| Uterus, NOS | 0.0 | 0.0 | 4.9 | 0 | ^ | ^ | ^ | ^ |
| Ovary | 0.0 | 0.0 | 4.9 | 0 | 8.2 | 5.0 | 13.0 | 22 |
| Vagina | 0.0 | 0.0 | 4.9 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 4.9 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 4.9 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 90.9 | 76.3 | 108.3 | 177 | 0.0 | 0.0 | 2.4 | 0 |
| Prostate | 83.9 | 69.7 | 100.8 | 154 | 0.0 | 0.0 | 2.4 | 0 |
| Testis | 5.0 | 3.0 | 10.1 | 20 | 0.0 | 0.0 | 2.4 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Other Male Genital Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Urinary System | 50.0 | 39.6 | 63.2 | 102 | 18.9 | 13.6 | 25.8 | 45 |
| Urinary Bladder | 19.0 | 12.6 | 28.5 | 36 | 5.2 | 2.5 | 9.9 | 11 |
| Kidney and Renal Pelvis | 29.5 | 21.8 | 40.0 | 63 | 13.2 | 8.9 | 19.1 | 33 |
| Ureter | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Urinary Organs | 0.0 | 0.0 | 4.9 | 0 | 0.0 | 0.0 | 2.4 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

| | Total | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Asian or Pacific Islander | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 13.8 | 0 |
| Pancreas | 17.7 | 7.5 | 40.6 | 9 | 9.1 | 2.5 | 26.6 | 6 |
| Retroperitoneum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 13.8 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 17.8 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 62.4 | 41.7 | 94.2 | 34 | 44.1 | 29.8 | 66.8 | 37 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 17.8 | 0 | ^ | ^ | ^ | ^ |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 13.8 | 0 |
| Lung and Bronchus | 56.1 | 36.6 | 86.8 | 31 | 42.4 | 28.2 | 64.9 | 35 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 13.8 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | 0.0 | 0.0 | 17.8 | 0 | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | 0.0 | 0.0 | 17.8 | 0 | 62.2 | 47.3 | 84.6 | 72 |
| Female Genital System | 0.0 | 0.0 | 17.8 | 0 | 34.8 | 24.6 | 53.0 | 40 |
| Cervix Uteri | 0.0 | 0.0 | 17.8 | 0 | 8.5 | 4.2 | 22.6 | 11 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 17.8 | 0 | 13.7 | 7.7 | 28.8 | 16 |
| Corpus Uteri | 0.0 | 0.0 | 17.8 | 0 | 13.7 | 7.7 | 28.8 | 16 |
| Uterus, NOS | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Ovary | 0.0 | 0.0 | 17.8 | 0 | 9.9 | 4.6 | 25.1 | 10 |
| Vagina | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Vulva | 0.0 | 0.0 | 17.8 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 17.8 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 141.9 | 106.3 | 188.6 | 66 | 0.0 | 0.0 | 13.8 | 0 |
| Prostate | 139.7 | 104.2 | 186.3 | 63 | 0.0 | 0.0 | 13.8 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 13.8 | 0 |
| Penis | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Urinary System | 32.8 | 17.1 | 60.8 | 15 | 14.1 | 5.8 | 32.5 | 10 |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 14.8 | 6.3 | 36.0 | 9 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Other Urinary Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 13.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

| | Total | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | White | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.8 | 0.4 | 1.9 | 14 | 0.4 | 0.1 | 1.1 | 6 |
| Brain and Other Nervous System | 7.8 | 6.4 | 9.8 | 134 | 6.4 | 5.1 | 8.1 | 95 |
| Brain | 7.5 | 6.2 | 9.5 | 129 | 6.2 | 5.0 | 7.8 | 92 |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Endocrine System | 4.9 | 3.9 | 6.6 | 89 | 12.5 | 10.8 | 14.4 | 222 |
| Thyroid | 4.3 | 3.3 | 5.8 | 79 | 11.9 | 10.3 | 13.8 | 213 |
| Other Endocrine including Thymus | 0.7 | 0.3 | 1.8 | 10 | 0.6 | 0.2 | 1.3 | 9 |
| Lymphoma | 30.4 | 26.9 | 34.5 | 408 | 20.8 | 18.2 | 23.7 | 272 |
| Hodgkin Lymphoma | 3.8 | 2.9 | 5.4 | 66 | 2.3 | 1.6 | 3.3 | 41 |
| Hodgkin - Nodal | 3.8 | 2.9 | 5.4 | 66 | 2.2 | 1.6 | 3.2 | 40 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 1.1 | 0 | ^ | ^ | ^ | ^ |
| Non-Hodgkin Lymphoma | 26.6 | 23.2 | 30.5 | 342 | 18.5 | 16.0 | 21.3 | 231 |
| NHL - Nodal | 18.4 | 15.7 | 21.6 | 247 | 12.6 | 10.6 | 14.9 | 160 |
| NHL - Extranodal | 8.2 | 6.3 | 10.8 | 95 | 5.9 | 4.5 | 7.6 | 71 |
| Myeloma | 5.3 | 3.9 | 7.2 | 64 | 4.6 | 3.3 | 6.2 | 51 |
| Leukemia | 17.2 | 14.5 | 20.5 | 208 | 11.2 | 9.2 | 13.5 | 132 |
| Lymphocytic Leukemia | 8.8 | 6.9 | 11.2 | 111 | 5.2 | 3.8 | 6.8 | 58 |
| Acute Lymphocytic Leukemia | 1.8 | 1.1 | 3.3 | 29 | 0.9 | 0.5 | 1.6 | 17 |
| Chronic Lymphocytic Leukemia | 6.4 | 4.7 | 8.6 | 71 | 4.0 | 2.8 | 5.5 | 38 |
| Other Lymphocytic Leukemia | 0.6 | 0.3 | 1.7 | 11 | ^ | ^ | ^ | ^ |
| Myeloid and Monocytic Leukemia | 7.2 | 5.4 | 9.6 | 82 | 4.9 | 3.6 | 6.5 | 60 |
| Acute Myeloid Leukemia | 4.1 | 2.9 | 6.0 | 50 | 2.6 | 1.7 | 3.9 | 33 |
| Acute Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Myeloid Leukemia | 2.9 | 1.7 | 4.8 | 28 | 1.7 | 1.0 | 2.8 | 20 |
| Other Myeloid/Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Leukemia | 1.2 | 0.6 | 2.5 | 15 | 1.1 | 0.6 | 2.1 | 14 |
| Other Acute Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Aleukemic, Subleukemic and NOS | 0.7 | 0.3 | 1.9 | 10 | 0.9 | 0.4 | 1.8 | 11 |
| Mesothelioma | 4.9 | 3.3 | 7.1 | 39 | ^ | ^ | ^ | ^ |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Miscellaneous | 11.5 | 9.2 | 14.4 | 126 | 9.6 | 7.7 | 11.8 | 101 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

| | Total | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Black | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 20.1 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 50.6 | 0 | ^ | ^ | ^ | ^ |
| Brain | 0.0 | 0.0 | 50.6 | 0 | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | 13.1 | 6.0 | 34.6 | 11 |
| Thyroid | ^ | ^ | ^ | ^ | 10.9 | 4.4 | 32.1 | 9 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphoma | 22.7 | 10.6 | 72.8 | 14 | 7.9 | 2.0 | 29.7 | 6 |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Non-Hodgkin Lymphoma | 21.4 | 9.5 | 71.6 | 12 | ^ | ^ | ^ | ^ |
| NHL - Nodal | 11.4 | 3.7 | 60.8 | 8 | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Leukemia | ^ | ^ | ^ | ^ | 9.5 | 3.0 | 31.4 | 6 |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 20.1 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Chronic Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 20.1 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | 9.5 | 3.0 | 31.4 | 6 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 50.6 | 0 | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Other Leukemia | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Mesothelioma | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 50.6 | 0 | 0.0 | 0.0 | 20.1 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

| | Total | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | American Indian/Alaska Native | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Brain and Other Nervous System | 3.6 | 1.6 | 9.0 | 12 | 1.8 | 0.7 | 4.5 | 8 |
| Brain | 3.4 | 1.4 | 8.8 | 11 | 1.8 | 0.7 | 4.5 | 8 |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Endocrine System | 2.1 | 0.8 | 7.0 | 7 | 11.4 | 8.0 | 16.3 | 40 |
| Thyroid | ^ | ^ | ^ | ^ | 11.4 | 8.0 | 16.3 | 40 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Lymphoma | 13.9 | 9.3 | 21.3 | 38 | 11.3 | 7.6 | 16.5 | 34 |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 4.9 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Non-Hodgkin Lymphoma | 12.9 | 8.5 | 20.2 | 35 | 10.5 | 7.0 | 15.6 | 31 |
| NHL - Nodal | 6.1 | 3.5 | 11.9 | 18 | 6.3 | 3.7 | 10.6 | 19 |
| NHL - Extranodal | 6.8 | 3.5 | 13.3 | 17 | 4.2 | 2.1 | 8.0 | 12 |
| Myeloma | ^ | ^ | ^ | ^ | 5.3 | 2.8 | 9.4 | 14 |
| Leukemia | 9.1 | 5.4 | 16.0 | 28 | 5.0 | 2.8 | 8.8 | 18 |
| Lymphocytic Leukemia | 1.8 | 0.8 | 6.5 | 10 | 2.7 | 1.2 | 6.1 | 10 |
| Acute Lymphocytic Leukemia | 1.4 | 0.6 | 6.1 | 9 | 1.4 | 0.5 | 4.0 | 7 |
| Chronic Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 4.9 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Myeloid and Monocytic Leukemia | 6.7 | 3.4 | 13.5 | 16 | 2.2 | 0.9 | 5.3 | 8 |
| Acute Myeloid Leukemia | 5.0 | 2.0 | 11.6 | 10 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 4.9 | 0 | 0.0 | 0.0 | 2.4 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Other Acute Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Mesothelioma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 2.4 | 0 |
| Miscellaneous | 15.5 | 9.7 | 24.4 | 27 | 12.8 | 8.3 | 19.1 | 29 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

| | Total | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Asian or Pacific Islander | | | | | | | |
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | 21.9 | 13.7 | 38.6 | 25 |
| Thyroid | ^ | ^ | ^ | ^ | 21.9 | 13.7 | 38.6 | 25 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Lymphoma | 9.0 | 3.6 | 27.6 | 8 | 8.9 | 3.6 | 24.2 | 8 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 17.8 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 0.0 | 0.0 | 17.8 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Non-Hodgkin Lymphoma | 9.0 | 3.6 | 27.6 | 8 | 8.3 | 3.2 | 23.5 | 7 |
| NHL - Nodal | 5.9 | 2.0 | 23.7 | 6 | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | 13.9 | 5.6 | 35.1 | 9 | ^ | ^ | ^ | ^ |
| Leukemia | 10.0 | 3.2 | 30.3 | 7 | 9.4 | 4.2 | 24.5 | 10 |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 17.8 | 0 | ^ | ^ | ^ | ^ |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | 5.9 | 1.9 | 20.5 | 6 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Other Leukemia | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Mesothelioma | 0.0 | 0.0 | 17.8 | 0 | 0.0 | 0.0 | 13.8 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 13.8 | 0 |
| Miscellaneous | 9.9 | 3.1 | 30.1 | 6 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1996

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 522.4 | 474.3 | 576.7 | 656 | 456.9 | 419.1 | 498.0 | 657 |
| Oral Cavity and Pharynx | 15.8 | 9.4 | 29.9 | 24 | 11.2 | 5.7 | 20.9 | 13 |
| Lip | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tongue | 4.0 | 1.6 | 16.2 | 7 | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.6 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nasopharynx | 0.0 | 0.0 | 12.3 | 0 | 0.0 | 0.0 | 5.6 | 0 |
| Tonsil | 4.8 | 1.4 | 17.9 | 7 | 0.0 | 0.0 | 5.6 | 0 |
| Oropharynx | 0.0 | 0.0 | 12.3 | 0 | ^ | ^ | ^ | ^ |
| Hypopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 12.3 | 0 | 0.0 | 0.0 | 5.6 | 0 |
| Digestive System | 87.3 | 67.5 | 114.0 | 104 | 77.7 | 61.3 | 97.9 | 91 |
| Esophagus | 5.5 | 0.9 | 20.5 | 6 | ^ | ^ | ^ | ^ |
| Stomach | 6.0 | 1.8 | 19.5 | 6 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 51.3 | 36.9 | 72.7 | 63 | 47.3 | 34.9 | 63.4 | 56 |
| Colon excluding Rectum | 33.2 | 21.7 | 52.1 | 39 | 34.8 | 24.0 | 49.5 | 38 |
| Cecum | 5.7 | 1.0 | 20.9 | 6 | 7.6 | 2.9 | 16.7 | 7 |
| Appendix | 0.0 | 0.0 | 12.3 | 0 | 0.0 | 0.0 | 5.6 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | 7.0 | 2.6 | 15.8 | 7 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.6 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 9.9 | 4.8 | 23.4 | 14 | 8.3 | 3.7 | 17.1 | 10 |
| Large Intestine, NOS | 6.4 | 2.1 | 20.1 | 6 | 4.8 | 1.7 | 12.1 | 7 |
| Rectum and Rectosigmoid Junction | 18.1 | 10.4 | 33.8 | 24 | 12.4 | 7.1 | 21.6 | 18 |
| Rectosigmoid Junction | 5.1 | 1.5 | 18.3 | 7 | ^ | ^ | ^ | ^ |
| Rectum | 13.0 | 6.7 | 27.7 | 17 | 8.8 | 4.6 | 16.7 | 14 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | 7.9 | 3.0 | 21.7 | 8 | 5.6 | 2.0 | 13.5 | 7 |
| Liver | 7.4 | 2.7 | 21.2 | 7 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.6 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1996

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 948.7 | 523.3 | 1,790.4 | 29 | 383.3 | 193.7 | 728.2 | 17 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 220.0 | 0 |
| Lip | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Tongue | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Salivary Gland | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 220.0 | 0 |
| Nasopharynx | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Tonsil | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Oropharynx | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Hypopharynx | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Digestive System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Esophagus | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 220.0 | 0 |
| Small Intestine | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 220.0 | 0 |
| Appendix | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Ascending Colon | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Transverse Colon | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Descending Colon | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 220.0 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 220.0 | 0 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 220.0 | 0 |
| Rectum | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Liver and Intrahepatic Bile Duct | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Liver | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Gallbladder | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1996

| | American Indian/Alaska Native | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 526.6 | 425.9 | 649.8 | 119 | 444.6 | 366.8 | 536.9 | 126 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Tongue | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.9 | 0 |
| Tonsil | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |
| Oropharynx | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.9 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Digestive System | 166.7 | 109.2 | 249.6 | 35 | 92.8 | 59.2 | 142.0 | 26 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | 27.6 | 12.4 | 69.2 | 9 | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Colon and Rectum | 114.1 | 65.3 | 190.0 | 21 | 64.3 | 37.0 | 107.8 | 18 |
| Colon excluding Rectum | 96.4 | 51.4 | 169.5 | 17 | 42.9 | 21.4 | 81.0 | 12 |
| Cecum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.9 | 0 |
| Sigmoid Colon | 47.5 | 19.7 | 103.4 | 10 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | 21.4 | 7.3 | 53.5 | 6 |
| Rectosigmoid Junction | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |
| Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Liver and Intrahepatic Bile Duct | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |
| Liver | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Gallbladder | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1996

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 493.5 | 301.8 | 811.8 | 25 | 300.1 | 157.4 | 589.9 | 22 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Lip | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Tongue | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Salivary Gland | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Nasopharynx | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Tonsil | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Oropharynx | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Hypopharynx | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Digestive System | 199.8 | 95.1 | 440.4 | 11 | 108.3 | 37.0 | 345.9 | 7 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 197.3 | 0 | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Ascending Colon | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | 0.0 | 0.0 | 197.3 | 0 | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Descending Colon | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Sigmoid Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 220.0 | 0 |
| Large Intestine, NOS | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Rectosigmoid Junction | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Rectum | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Gallbladder | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1996

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 12.3 | 0 | ^ | ^ | ^ | ^ |
| Pancreas | 12.5 | 5.3 | 28.6 | 13 | 13.1 | 6.9 | 23.5 | 15 |
| Retroperitoneum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 12.3 | 0 | 0.0 | 0.0 | 5.6 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 12.3 | 0 | ^ | ^ | ^ | ^ |
| Respiratory System | 99.5 | 79.2 | 126.6 | 121 | 71.7 | 56.7 | 90.3 | 88 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 12.3 | 0 | 0.0 | 0.0 | 5.6 | 0 |
| Larynx | 12.5 | 6.4 | 26.9 | 15 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 87.1 | 67.9 | 113.0 | 106 | 69.3 | 54.5 | 87.7 | 84 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 12.3 | 0 | 0.0 | 0.0 | 5.6 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | 3.4 | 1.4 | 15.3 | 8 | 3.2 | 1.2 | 9.2 | 7 |
| Skin excluding Basal and Squamous | 16.1 | 9.4 | 30.7 | 25 | 12.6 | 7.6 | 21.3 | 23 |
| Melanoma of the Skin | 16.1 | 9.4 | 30.7 | 25 | 12.6 | 7.6 | 21.3 | 23 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 12.3 | 0 | 0.0 | 0.0 | 5.6 | 0 |
| Breast | ^ | ^ | ^ | ^ | 132.4 | 113.3 | 154.7 | 210 |
| Female Genital System | 0.0 | 0.0 | 12.3 | 0 | 47.4 | 36.7 | 61.3 | 80 |
| Cervix Uteri | 0.0 | 0.0 | 12.3 | 0 | 11.3 | 7.0 | 19.1 | 23 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 12.3 | 0 | 21.5 | 14.2 | 32.4 | 33 |
| Corpus Uteri | 0.0 | 0.0 | 12.3 | 0 | 18.1 | 11.6 | 28.3 | 28 |
| Uterus, NOS | 0.0 | 0.0 | 12.3 | 0 | ^ | ^ | ^ | ^ |
| Ovary | 0.0 | 0.0 | 12.3 | 0 | 12.5 | 7.3 | 21.3 | 21 |
| Vagina | 0.0 | 0.0 | 12.3 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 12.3 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 12.3 | 0 | 0.0 | 0.0 | 5.6 | 0 |
| Male Genital System | 163.1 | 135.5 | 197.4 | 187 | 0.0 | 0.0 | 5.6 | 0 |
| Prostate | 158.0 | 130.5 | 192.2 | 172 | 0.0 | 0.0 | 5.6 | 0 |
| Testis | 5.1 | 2.8 | 16.8 | 15 | 0.0 | 0.0 | 5.6 | 0 |
| Penis | 0.0 | 0.0 | 12.3 | 0 | 0.0 | 0.0 | 5.6 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 12.3 | 0 | 0.0 | 0.0 | 5.6 | 0 |
| Urinary System | 57.0 | 41.8 | 79.3 | 69 | 17.5 | 10.5 | 28.4 | 22 |
| Urinary Bladder | 36.2 | 23.4 | 56.5 | 40 | 9.9 | 4.9 | 18.9 | 12 |
| Kidney and Renal Pelvis | 17.4 | 10.7 | 31.9 | 25 | 6.7 | 2.7 | 14.9 | 9 |
| Ureter | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Urinary Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.6 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1996

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Pancreas | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Respiratory System | 178.5 | 60.6 | 806.2 | 9 | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Larynx | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Lung and Bronchus | 160.0 | 48.4 | 789.2 | 8 | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 220.0 | 0 |
| Bones and Joints | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Breast | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Female Genital System | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Cervix Uteri | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Corpus Uteri | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Uterus, NOS | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Ovary | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Vagina | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Vulva | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Male Genital System | 392.4 | 175.1 | 1,066.4 | 11 | 0.0 | 0.0 | 220.0 | 0 |
| Prostate | 392.4 | 175.1 | 1,066.4 | 11 | 0.0 | 0.0 | 220.0 | 0 |
| Testis | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Penis | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 220.0 | 0 |
| Ureter | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1996

| | American Indian/Alaska Native | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Respiratory System | 137.4 | 91.4 | 206.9 | 33 | 70.3 | 40.5 | 116.5 | 18 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.9 | 0 |
| Larynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 122.6 | 78.8 | 190.2 | 29 | 66.0 | 37.3 | 111.2 | 17 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.9 | 0 |
| Bones and Joints | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Breast | 0.0 | 0.0 | 37.4 | 0 | 159.8 | 114.7 | 220.1 | 46 |
| Female Genital System | 0.0 | 0.0 | 37.4 | 0 | 35.3 | 17.1 | 69.7 | 11 |
| Cervix Uteri | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Ovary | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Vulva | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Male Genital System | 109.4 | 66.2 | 177.6 | 23 | 0.0 | 0.0 | 22.9 | 0 |
| Prostate | 101.8 | 59.6 | 169.5 | 20 | 0.0 | 0.0 | 22.9 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.9 | 0 |
| Penis | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Urinary System | 50.9 | 22.8 | 106.3 | 10 | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.9 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1996

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 220.0 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Larynx | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Bones and Joints | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 220.0 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Breast | 0.0 | 0.0 | 197.3 | 0 | ^ | ^ | ^ | ^ |
| Female Genital System | 0.0 | 0.0 | 197.3 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 197.3 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 197.3 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Ovary | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Vagina | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Vulva | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Male Genital System | 151.3 | 49.8 | 404.0 | 7 | 0.0 | 0.0 | 220.0 | 0 |
| Prostate | 151.3 | 49.8 | 404.0 | 7 | 0.0 | 0.0 | 220.0 | 0 |
| Testis | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Penis | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Urinary System | 0.0 | 0.0 | 197.3 | 0 | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 197.3 | 0 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Ureter | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1996

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain and Other Nervous System | 6.9 | 3.7 | 19.1 | 15 | 4.3 | 1.9 | 10.6 | 9 |
| Brain | 6.6 | 3.4 | 18.8 | 14 | 4.3 | 1.9 | 10.6 | 9 |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.6 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | 17.2 | 11.4 | 26.4 | 33 |
| Thyroid | ^ | ^ | ^ | ^ | 16.7 | 11.0 | 25.9 | 32 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphoma | 23.6 | 15.9 | 38.6 | 41 | 24.5 | 16.3 | 36.4 | 37 |
| Hodgkin Lymphoma | 5.6 | 2.7 | 17.7 | 12 | 4.9 | 2.5 | 11.1 | 12 |
| Hodgkin - Nodal | 5.6 | 2.7 | 17.7 | 12 | 4.4 | 2.2 | 10.5 | 11 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 12.3 | 0 | ^ | ^ | ^ | ^ |
| Non-Hodgkin Lymphoma | 18.0 | 11.1 | 32.6 | 29 | 19.6 | 12.0 | 31.1 | 25 |
| NHL - Nodal | 13.2 | 7.8 | 26.7 | 23 | 13.8 | 7.8 | 23.7 | 19 |
| NHL - Extranodal | 4.7 | 1.3 | 17.9 | 6 | 5.8 | 1.9 | 14.3 | 6 |
| Myeloma | ^ | ^ | ^ | ^ | 8.1 | 3.9 | 16.4 | 11 |
| Leukemia | 24.4 | 15.3 | 41.2 | 31 | 15.4 | 8.6 | 26.3 | 18 |
| Lymphocytic Leukemia | 14.5 | 7.9 | 29.2 | 19 | 5.8 | 2.1 | 13.9 | 7 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 11.1 | 5.1 | 25.7 | 12 | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.6 | 0 |
| Myeloid and Monocytic Leukemia | 6.2 | 2.2 | 19.5 | 8 | 8.8 | 3.9 | 18.0 | 10 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 12.3 | 0 | ^ | ^ | ^ | ^ |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 12.3 | 0 | 0.0 | 0.0 | 5.6 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.6 | 0 |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Mesothelioma | 8.9 | 3.7 | 22.9 | 9 | 0.0 | 0.0 | 5.6 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 12.3 | 0 | 0.0 | 0.0 | 5.6 | 0 |
| Miscellaneous | 5.4 | 1.7 | 18.6 | 8 | 12.1 | 6.0 | 22.3 | 12 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1996

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Brain | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Endocrine System | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Thyroid | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Leukemia | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Mesothelioma | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Miscellaneous | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 220.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1996

| | American Indian/Alaska Native | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.9 | 0 |
| Brain | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.9 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.9 | 0 |
| Leukemia | 23.6 | 6.2 | 70.8 | 6 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.9 | 0 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.9 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Other Leukemia | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 37.4 | 0 | 0.0 | 0.0 | 22.9 | 0 |
| Mesothelioma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.9 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.9 | 0 |
| Miscellaneous | 0.0 | 0.0 | 37.4 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1996

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 197.3 | 0 | ^ | ^ | ^ | ^ |
| Brain | 0.0 | 0.0 | 197.3 | 0 | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Lymphoma | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Non-Hodgkin Lymphoma | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| NHL - Nodal | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| NHL - Extranodal | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 220.0 | 0 |
| Leukemia | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Lymphocytic Leukemia | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Leukemia | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Mesothelioma | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 197.3 | 0 | 0.0 | 0.0 | 220.0 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1997

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 593.0 | 542.5 | 649.1 | 759 | 431.4 | 395.6 | 470.2 | 638 |
| Oral Cavity and Pharynx | 13.5 | 7.9 | 26.4 | 23 | 7.0 | 3.2 | 14.5 | 11 |
| Lip | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tongue | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.1 | 0 |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.1 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Oropharynx | 0.0 | 0.0 | 11.3 | 0 | 0.0 | 0.0 | 5.1 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.1 | 0 |
| Digestive System | 91.7 | 72.9 | 116.7 | 118 | 57.6 | 44.4 | 74.1 | 76 |
| Esophagus | 9.8 | 4.3 | 23.2 | 12 | ^ | ^ | ^ | ^ |
| Stomach | 7.0 | 2.7 | 19.7 | 8 | 4.4 | 1.4 | 11.4 | 6 |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 46.8 | 34.2 | 65.7 | 61 | 31.6 | 21.9 | 44.8 | 41 |
| Colon excluding Rectum | 32.3 | 21.8 | 49.6 | 42 | 24.5 | 16.0 | 36.6 | 31 |
| Cecum | 4.7 | 1.6 | 16.6 | 7 | 7.6 | 3.1 | 16.3 | 8 |
| Appendix | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ascending Colon | 6.8 | 2.5 | 19.5 | 8 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.1 | 0 |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 11.6 | 5.3 | 25.7 | 12 | 6.2 | 2.5 | 14.0 | 8 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 14.4 | 8.3 | 28.0 | 19 | 7.1 | 3.2 | 14.8 | 10 |
| Rectosigmoid Junction | 5.1 | 1.8 | 17.1 | 6 | ^ | ^ | ^ | ^ |
| Rectum | 9.3 | 4.6 | 21.8 | 13 | 6.7 | 2.9 | 14.4 | 9 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | 5.6 | 2.2 | 12.9 | 8 |
| Liver and Intrahepatic Bile Duct | 9.7 | 3.9 | 23.6 | 14 | ^ | ^ | ^ | ^ |
| Liver | 5.6 | 2.6 | 16.9 | 11 | 0.0 | 0.0 | 5.1 | 0 |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1997

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 653.1 | 403.5 | 1,203.0 | 28 | 462.0 | 260.4 | 802.4 | 21 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 476.7 | 0 | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Tongue | 0.0 | 0.0 | 476.7 | 0 | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Nasopharynx | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Tonsil | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Oropharynx | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Hypopharynx | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Digestive System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Esophagus | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 204.3 | 0 |
| Small Intestine | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 204.3 | 0 |
| Appendix | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Ascending Colon | 0.0 | 0.0 | 476.7 | 0 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Transverse Colon | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Descending Colon | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 476.7 | 0 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 204.3 | 0 |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 476.7 | 0 | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Rectum | 0.0 | 0.0 | 476.7 | 0 | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Liver and Intrahepatic Bile Duct | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Liver | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Gallbladder | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1997

| | American Indian/Alaska Native | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 654.1 | 542.9 | 786.8 | 154 | 451.2 | 374.0 | 542.4 | 133 |
| Oral Cavity and Pharynx | 31.8 | 11.5 | 79.3 | 7 | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Tongue | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Salivary Gland | 0.0 | 0.0 | 36.0 | 0 | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Oropharynx | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Hypopharynx | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Digestive System | 274.0 | 202.6 | 368.2 | 61 | 172.1 | 123.7 | 235.7 | 46 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | 28.5 | 12.8 | 69.6 | 9 | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 36.0 | 0 | ^ | ^ | ^ | ^ |
| Colon and Rectum | 166.6 | 112.6 | 244.0 | 37 | 93.2 | 58.8 | 143.1 | 25 |
| Colon excluding Rectum | 120.8 | 73.7 | 192.8 | 24 | 58.6 | 31.5 | 102.2 | 15 |
| Cecum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Appendix | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | 0.0 | 0.0 | 36.0 | 0 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 52.0 | 21.8 | 109.9 | 9 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 45.8 | 23.2 | 92.8 | 13 | 34.5 | 16.1 | 69.0 | 10 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 36.1 | 16.2 | 81.3 | 10 | 28.3 | 11.8 | 61.4 | 8 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1997

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 529.8 | 344.6 | 817.9 | 34 | 271.0 | 181.7 | 470.0 | 32 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Lip | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Tongue | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Salivary Gland | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Oropharynx | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Hypopharynx | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Digestive System | 80.1 | 30.7 | 251.4 | 7 | 69.6 | 25.9 | 250.6 | 7 |
| Esophagus | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Stomach | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Ascending Colon | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Descending Colon | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Gallbladder | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1997

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.1 | 0 |
| Pancreas | 13.5 | 6.4 | 28.4 | 16 | 8.0 | 3.6 | 16.2 | 10 |
| Retroperitoneum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 11.3 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | 0.0 | 0.0 | 11.3 | 0 | ^ | ^ | ^ | ^ |
| Respiratory System | 107.4 | 86.6 | 134.4 | 129 | 56.6 | 43.6 | 72.9 | 73 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 11.3 | 0 | ^ | ^ | ^ | ^ |
| Larynx | 11.9 | 5.4 | 26.4 | 15 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 95.5 | 76.1 | 121.1 | 114 | 55.6 | 42.7 | 71.9 | 71 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 11.3 | 0 | 0.0 | 0.0 | 5.1 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 19.7 | 11.4 | 35.2 | 32 | 13.7 | 9.0 | 21.5 | 29 |
| Melanoma of the Skin | 18.8 | 10.7 | 34.2 | 30 | 13.3 | 8.6 | 21.0 | 28 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 149.2 | 129.0 | 172.3 | 233 |
| Female Genital System | 0.0 | 0.0 | 11.3 | 0 | 57.9 | 45.1 | 73.8 | 84 |
| Cervix Uteri | 0.0 | 0.0 | 11.3 | 0 | 11.0 | 6.8 | 18.4 | 23 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 11.3 | 0 | 25.8 | 17.1 | 37.9 | 33 |
| Corpus Uteri | 0.0 | 0.0 | 11.3 | 0 | 23.2 | 15.1 | 34.8 | 30 |
| Uterus, NOS | 0.0 | 0.0 | 11.3 | 0 | ^ | ^ | ^ | ^ |
| Ovary | 0.0 | 0.0 | 11.3 | 0 | 17.5 | 10.6 | 28.1 | 22 |
| Vagina | 0.0 | 0.0 | 11.3 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 11.3 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 11.3 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 187.9 | 160.1 | 221.7 | 222 | 0.0 | 0.0 | 5.1 | 0 |
| Prostate | 181.6 | 153.9 | 215.3 | 205 | 0.0 | 0.0 | 5.1 | 0 |
| Testis | 6.3 | 3.7 | 17.2 | 17 | 0.0 | 0.0 | 5.1 | 0 |
| Penis | 0.0 | 0.0 | 11.3 | 0 | 0.0 | 0.0 | 5.1 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 11.3 | 0 | 0.0 | 0.0 | 5.1 | 0 |
| Urinary System | 85.3 | 64.6 | 112.7 | 90 | 25.8 | 17.7 | 37.2 | 37 |
| Urinary Bladder | 67.8 | 48.6 | 94.2 | 63 | 14.6 | 8.5 | 24.4 | 19 |
| Kidney and Renal Pelvis | 15.2 | 9.3 | 28.2 | 25 | 11.1 | 6.3 | 19.4 | 18 |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.1 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 11.3 | 0 | 0.0 | 0.0 | 5.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1997

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Pancreas | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Larynx | 0.0 | 0.0 | 476.7 | 0 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Bones and Joints | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Breast | 0.0 | 0.0 | 476.7 | 0 | 133.3 | 37.4 | 390.2 | 6 |
| Female Genital System | 0.0 | 0.0 | 476.7 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 476.7 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Corpus Uteri | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Uterus, NOS | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Ovary | 0.0 | 0.0 | 476.7 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Vulva | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Male Genital System | 272.6 | 132.7 | 756.6 | 12 | 0.0 | 0.0 | 204.3 | 0 |
| Prostate | 266.9 | 127.8 | 751.3 | 11 | 0.0 | 0.0 | 204.3 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 204.3 | 0 |
| Penis | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 204.3 | 0 |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1997

| | American Indian/Alaska Native | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 36.0 | 0 | ^ | ^ | ^ | ^ |
| Pancreas | ^ | ^ | ^ | ^ | 31.1 | 12.1 | 67.8 | 7 |
| Retroperitoneum | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Respiratory System | 165.4 | 112.6 | 241.3 | 39 | 68.4 | 39.7 | 113.0 | 18 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Larynx | 0.0 | 0.0 | 36.0 | 0 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 153.0 | 102.5 | 226.7 | 36 | 64.3 | 36.6 | 108.0 | 17 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 36.0 | 0 | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | 0.0 | 0.0 | 36.0 | 0 | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Breast | 0.0 | 0.0 | 36.0 | 0 | 114.2 | 79.4 | 163.1 | 37 |
| Female Genital System | 0.0 | 0.0 | 36.0 | 0 | 25.6 | 10.9 | 56.4 | 8 |
| Cervix Uteri | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 36.0 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 36.0 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Ovary | 0.0 | 0.0 | 36.0 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Vulva | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Male Genital System | 93.2 | 50.7 | 162.0 | 17 | 0.0 | 0.0 | 22.0 | 0 |
| Prostate | 91.2 | 49.0 | 159.9 | 16 | 0.0 | 0.0 | 22.0 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Penis | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Urinary System | 36.8 | 16.6 | 82.3 | 10 | 26.3 | 9.2 | 61.4 | 6 |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 24.2 | 8.0 | 67.7 | 6 | ^ | ^ | ^ | ^ |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1997

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Respiratory System | 98.6 | 39.9 | 276.3 | 8 | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Larynx | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Lung and Bronchus | 98.6 | 39.9 | 276.3 | 8 | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Bones and Joints | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Breast | 0.0 | 0.0 | 163.5 | 0 | 73.1 | 34.4 | 247.8 | 10 |
| Female Genital System | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Ovary | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Vulva | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 221.0 | 94.6 | 469.7 | 10 | 0.0 | 0.0 | 178.8 | 0 |
| Prostate | 208.5 | 84.4 | 457.3 | 8 | 0.0 | 0.0 | 178.8 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Penis | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Ureter | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1997

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain and Other Nervous System | 6.4 | 2.7 | 18.3 | 13 | 7.5 | 3.8 | 14.7 | 13 |
| Brain | 6.1 | 2.5 | 18.0 | 12 | 7.5 | 3.8 | 14.7 | 13 |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.1 | 0 |
| Endocrine System | 3.5 | 1.6 | 14.4 | 9 | 11.0 | 6.6 | 18.7 | 23 |
| Thyroid | 3.5 | 1.6 | 14.4 | 9 | 11.0 | 6.6 | 18.7 | 23 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 11.3 | 0 | 0.0 | 0.0 | 5.1 | 0 |
| Lymphoma | 37.2 | 27.1 | 53.6 | 64 | 17.8 | 10.7 | 28.6 | 23 |
| Hodgkin Lymphoma | 4.1 | 1.8 | 15.2 | 9 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 4.1 | 1.8 | 15.2 | 9 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 11.3 | 0 | 0.0 | 0.0 | 5.1 | 0 |
| Non-Hodgkin Lymphoma | 33.1 | 23.4 | 49.3 | 55 | 16.3 | 9.4 | 26.8 | 20 |
| NHL - Nodal | 23.0 | 15.1 | 37.8 | 38 | 10.9 | 5.4 | 20.1 | 14 |
| NHL - Extranodal | 10.1 | 5.2 | 22.8 | 17 | 5.4 | 1.9 | 12.9 | 6 |
| Myeloma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Leukemia | 18.2 | 10.5 | 33.3 | 28 | 10.4 | 5.2 | 19.4 | 14 |
| Lymphocytic Leukemia | 11.4 | 5.0 | 26.0 | 15 | 4.6 | 1.4 | 11.8 | 6 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 9.4 | 3.4 | 23.9 | 10 | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 11.3 | 0 | ^ | ^ | ^ | ^ |
| Myeloid and Monocytic Leukemia | 5.3 | 2.2 | 16.8 | 10 | 4.4 | 1.4 | 11.4 | 6 |
| Acute Myeloid Leukemia | 3.8 | 1.2 | 15.4 | 6 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 11.3 | 0 | 0.0 | 0.0 | 5.1 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 11.3 | 0 | 0.0 | 0.0 | 5.1 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | 0.0 | 0.0 | 11.3 | 0 | ^ | ^ | ^ | ^ |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Mesothelioma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.1 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 5.1 | 0 |
| Miscellaneous | 9.1 | 3.3 | 23.3 | 11 | 9.0 | 4.2 | 17.7 | 10 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1997

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Brain | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 204.3 | 0 |
| Thyroid | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 204.3 | 0 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 204.3 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 204.3 | 0 |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 204.3 | 0 |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 204.3 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 204.3 | 0 |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 204.3 | 0 |
| NHL - Extranodal | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Myeloma | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Leukemia | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Lymphocytic Leukemia | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Other Leukemia | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Mesothelioma | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |
| Miscellaneous | 0.0 | 0.0 | 476.7 | 0 | 0.0 | 0.0 | 204.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1997

| | American Indian/Alaska Native | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Brain | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 36.0 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Lymphoma | 16.4 | 5.9 | 54.1 | 6 | 14.1 | 4.9 | 39.8 | 6 |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 14.1 | 4.9 | 39.8 | 6 |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Chronic Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 36.0 | 0 | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Other Acute Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Mesothelioma | 0.0 | 0.0 | 36.0 | 0 | 0.0 | 0.0 | 22.0 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 22.0 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1997

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Brain | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Endocrine System | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Leukemia | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 163.5 | 0 | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Leukemia | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Mesothelioma | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Miscellaneous | 0.0 | 0.0 | 163.5 | 0 | 0.0 | 0.0 | 178.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1998

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 563.4 | 516.8 | 615.1 | 753 | 473.1 | 436.7 | 512.5 | 734 |
| Oral Cavity and Pharynx | 9.1 | 5.3 | 19.6 | 20 | 8.9 | 4.6 | 16.7 | 14 |
| Lip | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tongue | 2.9 | 1.0 | 12.8 | 6 | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 10.1 | 0 | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Oropharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.8 | 0 |
| Hypopharynx | 0.0 | 0.0 | 10.1 | 0 | 0.0 | 0.0 | 4.8 | 0 |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.8 | 0 |
| Digestive System | 104.8 | 84.2 | 131.0 | 131 | 73.5 | 58.6 | 91.5 | 96 |
| Esophagus | 5.0 | 2.0 | 15.4 | 8 | ^ | ^ | ^ | ^ |
| Stomach | 15.6 | 7.3 | 31.2 | 13 | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 10.1 | 0 | ^ | ^ | ^ | ^ |
| Colon and Rectum | 53.8 | 39.2 | 74.5 | 65 | 47.1 | 35.3 | 62.2 | 60 |
| Colon excluding Rectum | 39.4 | 26.1 | 59.1 | 41 | 32.7 | 22.9 | 45.9 | 39 |
| Cecum | 8.9 | 2.7 | 23.1 | 8 | 13.9 | 7.8 | 23.6 | 16 |
| Appendix | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.8 | 0 |
| Ascending Colon | 11.1 | 4.0 | 26.1 | 8 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 10.1 | 0 | 0.0 | 0.0 | 4.8 | 0 |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 8.6 | 4.1 | 20.0 | 11 | 10.7 | 5.3 | 19.7 | 12 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 14.4 | 9.0 | 26.2 | 24 | 14.4 | 8.5 | 23.7 | 21 |
| Rectosigmoid Junction | 5.0 | 1.8 | 15.7 | 6 | ^ | ^ | ^ | ^ |
| Rectum | 9.4 | 5.4 | 20.2 | 18 | 11.3 | 6.1 | 20.0 | 16 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | 6.3 | 3.1 | 16.7 | 12 | 5.2 | 1.7 | 12.7 | 6 |
| Liver | 6.3 | 3.1 | 16.7 | 12 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 10.1 | 0 | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1998

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 753.3 | 464.8 | 1,449.2 | 28 | 626.5 | 375.4 | 1,029.1 | 28 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| Lip | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Tongue | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| Salivary Gland | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Nasopharynx | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Tonsil | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Oropharynx | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Hypopharynx | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Digestive System | 260.5 | 102.7 | 902.0 | 9 | ^ | ^ | ^ | ^ |
| Esophagus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| Small Intestine | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| Appendix | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Ascending Colon | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Descending Colon | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| Large Intestine, NOS | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Rectum | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| Liver | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1998

| | American Indian/Alaska Native | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 529.0 | 430.9 | 648.2 | 132 | 528.9 | 446.4 | 624.7 | 164 |
| Oral Cavity and Pharynx | 46.1 | 21.9 | 94.9 | 13 | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Tongue | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Salivary Gland | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Oropharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Hypopharynx | 0.0 | 0.0 | 34.9 | 0 | ^ | ^ | ^ | ^ |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Digestive System | 166.9 | 114.2 | 242.1 | 42 | 208.4 | 155.0 | 276.6 | 56 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | 23.6 | 9.7 | 62.7 | 8 | 17.1 | 5.7 | 44.6 | 6 |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 92.5 | 53.6 | 155.5 | 21 | 141.0 | 96.6 | 200.6 | 36 |
| Colon excluding Rectum | 74.6 | 39.1 | 135.2 | 16 | 109.1 | 70.4 | 163.5 | 27 |
| Cecum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Sigmoid Colon | 26.2 | 8.0 | 71.7 | 6 | 39.4 | 18.8 | 75.9 | 11 |
| Large Intestine, NOS | 0.0 | 0.0 | 34.9 | 0 | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | 31.8 | 13.5 | 66.5 | 9 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | ^ | ^ | ^ | ^ | 20.1 | 6.5 | 50.5 | 6 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 34.9 | 0 | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1998

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 567.1 | 359.7 | 887.2 | 30 | 210.5 | 132.4 | 392.0 | 24 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 163.5 | 0 |
| Lip | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Tongue | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Salivary Gland | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 163.5 | 0 |
| Nasopharynx | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Tonsil | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Oropharynx | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Hypopharynx | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Digestive System | 79.9 | 30.3 | 259.9 | 7 | 53.8 | 19.2 | 216.7 | 6 |
| Esophagus | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Stomach | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 163.5 | 0 |
| Cecum | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Appendix | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Ascending Colon | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Transverse Colon | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Descending Colon | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 163.5 | 0 |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 173.4 | 0 | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Rectum | 0.0 | 0.0 | 173.4 | 0 | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 163.5 | 0 |
| Gallbladder | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1998

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 19.9 | 11.9 | 34.3 | 26 | 8.2 | 4.1 | 15.7 | 13 |
| Retroperitoneum | 0.0 | 0.0 | 10.1 | 0 | 0.0 | 0.0 | 4.8 | 0 |
| Peritoneum, Omentum and Mesentery | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.8 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 93.6 | 75.6 | 116.9 | 125 | 79.2 | 63.9 | 97.5 | 103 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Larynx | 4.1 | 1.6 | 14.4 | 8 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 88.7 | 71.1 | 111.7 | 116 | 77.3 | 62.2 | 95.5 | 100 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 10.1 | 0 | 0.0 | 0.0 | 4.8 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 3.9 | 1.6 | 9.7 | 8 |
| Skin excluding Basal and Squamous | 16.5 | 10.6 | 28.5 | 30 | 9.6 | 5.8 | 16.5 | 21 |
| Melanoma of the Skin | 14.2 | 8.8 | 25.9 | 26 | 6.7 | 3.8 | 12.8 | 16 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 156.7 | 136.5 | 179.7 | 261 |
| Female Genital System | 0.0 | 0.0 | 10.1 | 0 | 52.6 | 41.2 | 66.9 | 87 |
| Cervix Uteri | 0.0 | 0.0 | 10.1 | 0 | 13.2 | 8.0 | 21.5 | 25 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 10.1 | 0 | 25.1 | 17.4 | 35.9 | 39 |
| Corpus Uteri | 0.0 | 0.0 | 10.1 | 0 | 25.1 | 17.4 | 35.9 | 39 |
| Uterus, NOS | 0.0 | 0.0 | 10.1 | 0 | 0.0 | 0.0 | 4.8 | 0 |
| Ovary | 0.0 | 0.0 | 10.1 | 0 | 10.1 | 5.6 | 17.9 | 18 |
| Vagina | 0.0 | 0.0 | 10.1 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 10.1 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 10.1 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 180.7 | 154.3 | 212.4 | 225 | 0.0 | 0.0 | 4.8 | 0 |
| Prostate | 172.1 | 146.4 | 203.1 | 209 | 0.0 | 0.0 | 4.8 | 0 |
| Testis | 3.9 | 1.9 | 13.6 | 11 | 0.0 | 0.0 | 4.8 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.8 | 0 |
| Other Male Genital Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.8 | 0 |
| Urinary System | 63.2 | 48.0 | 84.1 | 77 | 16.5 | 10.3 | 25.9 | 25 |
| Urinary Bladder | 46.6 | 33.1 | 66.2 | 49 | 10.2 | 5.3 | 18.5 | 14 |
| Kidney and Renal Pelvis | 14.8 | 9.1 | 26.8 | 26 | 6.3 | 3.0 | 13.0 | 11 |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.8 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 10.1 | 0 | 0.0 | 0.0 | 4.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1998

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Bones and Joints | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Breast | ^ | ^ | ^ | ^ | 259.1 | 119.4 | 557.9 | 15 |
| Female Genital System | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Corpus Uteri | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Uterus, NOS | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Ovary | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Vulva | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Male Genital System | 222.2 | 88.2 | 850.6 | 9 | 0.0 | 0.0 | 227.0 | 0 |
| Prostate | 222.2 | 88.2 | 850.6 | 9 | 0.0 | 0.0 | 227.0 | 0 |
| Testis | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Penis | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 635.7 | 0 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| Ureter | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1998

| | American Indian/Alaska Native | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 34.9 | 0 | ^ | ^ | ^ | ^ |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Peritoneum, Omentum and Mesentery | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Respiratory System | 140.4 | 93.8 | 209.3 | 35 | 69.0 | 40.5 | 112.7 | 19 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 34.9 | 0 | ^ | ^ | ^ | ^ |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Lung and Bronchus | 134.3 | 88.5 | 202.5 | 33 | 65.9 | 38.0 | 109.1 | 18 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Bones and Joints | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Breast | 0.0 | 0.0 | 34.9 | 0 | 158.2 | 115.6 | 214.5 | 50 |
| Female Genital System | 0.0 | 0.0 | 34.9 | 0 | 34.2 | 18.4 | 64.1 | 14 |
| Cervix Uteri | 0.0 | 0.0 | 34.9 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 34.9 | 0 | 15.5 | 5.6 | 41.2 | 6 |
| Corpus Uteri | 0.0 | 0.0 | 34.9 | 0 | 15.5 | 5.6 | 41.2 | 6 |
| Uterus, NOS | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Ovary | 0.0 | 0.0 | 34.9 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Vulva | 0.0 | 0.0 | 34.9 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Male Genital System | 89.0 | 49.6 | 153.0 | 18 | 0.0 | 0.0 | 21.1 | 0 |
| Prostate | 75.8 | 40.8 | 135.4 | 16 | 0.0 | 0.0 | 21.1 | 0 |
| Testis | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Urinary System | 36.3 | 15.2 | 82.6 | 10 | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Kidney and Renal Pelvis | 27.6 | 9.2 | 72.7 | 7 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1998

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Pancreas | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Retroperitoneum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 163.5 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 173.4 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 163.5 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Bones and Joints | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 163.5 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Breast | 0.0 | 0.0 | 173.4 | 0 | ^ | ^ | ^ | ^ |
| Female Genital System | 0.0 | 0.0 | 173.4 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 173.4 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 173.4 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 173.4 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Ovary | 0.0 | 0.0 | 173.4 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Vulva | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Male Genital System | 186.2 | 72.5 | 430.3 | 8 | 0.0 | 0.0 | 163.5 | 0 |
| Prostate | 186.2 | 72.5 | 430.3 | 8 | 0.0 | 0.0 | 163.5 | 0 |
| Testis | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Penis | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 163.5 | 0 |
| Kidney and Renal Pelvis | 0.0 | 0.0 | 173.4 | 0 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1998

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.8 | 0 |
| Brain and Other Nervous System | 9.0 | 5.1 | 19.6 | 19 | 6.4 | 3.1 | 13.0 | 12 |
| Brain | 9.0 | 5.1 | 19.6 | 19 | 6.4 | 3.1 | 13.0 | 12 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 10.1 | 0 | 0.0 | 0.0 | 4.8 | 0 |
| Endocrine System | 3.1 | 1.3 | 12.9 | 8 | 16.3 | 11.0 | 24.4 | 34 |
| Thyroid | 3.1 | 1.3 | 12.9 | 8 | 15.8 | 10.6 | 23.9 | 33 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 10.1 | 0 | ^ | ^ | ^ | ^ |
| Lymphoma | 32.6 | 22.0 | 49.2 | 48 | 23.5 | 16.1 | 34.0 | 38 |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 10.1 | 0 | 0.0 | 0.0 | 4.8 | 0 |
| Non-Hodgkin Lymphoma | 29.2 | 19.2 | 45.4 | 44 | 21.9 | 14.8 | 32.1 | 35 |
| NHL - Nodal | 20.2 | 12.4 | 34.3 | 32 | 10.5 | 5.9 | 18.4 | 17 |
| NHL - Extranodal | 9.1 | 3.6 | 21.8 | 12 | 11.4 | 6.4 | 19.7 | 18 |
| Myeloma | 9.9 | 5.3 | 21.3 | 15 | 6.7 | 2.9 | 14.0 | 9 |
| Leukemia | 24.0 | 14.3 | 40.3 | 29 | 7.7 | 3.5 | 15.4 | 11 |
| Lymphocytic Leukemia | 10.7 | 4.6 | 23.9 | 13 | 4.4 | 1.5 | 11.1 | 6 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 5.4 | 1.8 | 16.6 | 6 | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloid and Monocytic Leukemia | 11.3 | 4.9 | 25.0 | 13 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 6.2 | 2.1 | 17.7 | 7 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.8 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.8 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.8 | 0 |
| Other Acute Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.8 | 0 |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.8 | 0 |
| Mesothelioma | 6.2 | 2.1 | 17.8 | 6 | 0.0 | 0.0 | 4.8 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.8 | 0 |
| Miscellaneous | 5.9 | 2.5 | 16.6 | 9 | 11.3 | 6.0 | 20.1 | 14 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1998

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Brain | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Endocrine System | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Thyroid | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| NHL - Extranodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 227.0 | 0 |
| Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Lymphocytic Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Other Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Mesothelioma | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |
| Miscellaneous | 0.0 | 0.0 | 635.7 | 0 | 0.0 | 0.0 | 227.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1998

| | American Indian/Alaska Native | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Endocrine System | 0.0 | 0.0 | 34.9 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 34.9 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | 0.0 | 0.0 | 34.9 | 0 | ^ | ^ | ^ | ^ |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 34.9 | 0 | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 34.9 | 0 | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 34.9 | 0 | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Acute Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Other Leukemia | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 34.9 | 0 | 0.0 | 0.0 | 21.1 | 0 |
| Mesothelioma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 21.1 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1998

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | 0.0 | 0.0 | 173.4 | 0 | ^ | ^ | ^ | ^ |
| Myeloma | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 163.5 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 163.5 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 163.5 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Other Leukemia | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Mesothelioma | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |
| Miscellaneous | 0.0 | 0.0 | 173.4 | 0 | 0.0 | 0.0 | 163.5 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1999

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 554.4 | 510.4 | 602.9 | 805 | 453.2 | 417.8 | 491.4 | 708 |
| Oral Cavity and Pharynx | 14.0 | 8.0 | 25.6 | 26 | 7.0 | 2.9 | 14.7 | 8 |
| Lip | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.6 | 0 |
| Tongue | 6.9 | 2.5 | 17.9 | 11 | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Floor of Mouth | 0.0 | 0.0 | 8.9 | 0 | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | 0.0 | 0.0 | 8.9 | 0 | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Oropharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.6 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.6 | 0 |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.6 | 0 |
| Digestive System | 96.2 | 78.2 | 119.1 | 137 | 67.5 | 53.1 | 85.0 | 85 |
| Esophagus | 10.0 | 5.4 | 20.5 | 16 | ^ | ^ | ^ | ^ |
| Stomach | 8.6 | 3.0 | 21.3 | 10 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 59.9 | 45.9 | 79.1 | 80 | 39.1 | 28.2 | 53.2 | 47 |
| Colon excluding Rectum | 49.0 | 36.0 | 67.2 | 60 | 29.6 | 20.2 | 42.3 | 35 |
| Cecum | 8.0 | 3.6 | 18.6 | 12 | 11.5 | 6.0 | 20.5 | 13 |
| Appendix | 0.0 | 0.0 | 8.9 | 0 | 0.0 | 0.0 | 4.6 | 0 |
| Ascending Colon | 14.1 | 7.7 | 26.3 | 16 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | 4.5 | 1.5 | 14.2 | 7 | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 13.0 | 6.5 | 25.6 | 14 | 5.4 | 1.9 | 12.6 | 6 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 10.9 | 6.1 | 21.6 | 20 | 9.5 | 4.6 | 17.7 | 12 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 8.3 | 4.1 | 18.6 | 15 | 7.7 | 3.4 | 15.5 | 10 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 8.9 | 0 | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | 6.5 | 2.8 | 16.6 | 11 | ^ | ^ | ^ | ^ |
| Liver | 6.5 | 2.8 | 16.6 | 11 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 8.9 | 0 | 0.0 | 0.0 | 4.6 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1999

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 690.0 | 447.0 | 1,297.5 | 30 | 403.0 | 221.6 | 708.7 | 21 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Lip | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Tongue | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Salivary Gland | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Nasopharynx | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Tonsil | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Oropharynx | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Hypopharynx | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Digestive System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Esophagus | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Small Intestine | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Appendix | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Transverse Colon | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Descending Colon | 0.0 | 0.0 | 572.1 | 0 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Large Intestine, NOS | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Rectosigmoid Junction | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Rectum | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Liver | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Gallbladder | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1999

| | American Indian/Alaska Native | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 624.2 | 518.3 | 752.5 | 154 | 488.7 | 411.7 | 578.3 | 159 |
| Oral Cavity and Pharynx | 48.1 | 24.3 | 98.5 | 14 | ^ | ^ | ^ | ^ |
| Lip | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.6 | 0 |
| Tongue | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.6 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Nasopharynx | 18.2 | 5.9 | 60.6 | 6 | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.6 | 0 |
| Oropharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.6 | 0 |
| Hypopharynx | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Digestive System | 250.3 | 182.9 | 341.9 | 58 | 188.6 | 140.9 | 249.6 | 57 |
| Esophagus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.6 | 0 |
| Stomach | 25.4 | 9.6 | 70.0 | 7 | 33.2 | 15.6 | 65.3 | 11 |
| Small Intestine | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.6 | 0 |
| Colon and Rectum | 154.8 | 104.3 | 230.0 | 37 | 120.1 | 83.6 | 169.8 | 37 |
| Colon excluding Rectum | 114.7 | 73.3 | 180.8 | 29 | 91.0 | 59.8 | 135.5 | 28 |
| Cecum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | 20.3 | 7.1 | 48.9 | 6 |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | 0.0 | 0.0 | 39.5 | 0 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 55.5 | 30.8 | 105.9 | 16 | 23.4 | 10.1 | 50.9 | 8 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 40.2 | 15.3 | 94.7 | 8 | 29.1 | 12.7 | 60.2 | 9 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 34.2 | 11.1 | 88.2 | 6 | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.6 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1999

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 361.3 | 221.3 | 594.9 | 23 | 297.8 | 175.5 | 512.5 | 30 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Tongue | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Nasopharynx | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Tonsil | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Oropharynx | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Hypopharynx | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Digestive System | 135.8 | 55.6 | 322.2 | 9 | 86.6 | 23.3 | 262.9 | 7 |
| Esophagus | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Stomach | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 143.0 | 0 |
| Cecum | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Appendix | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Ascending Colon | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Transverse Colon | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Descending Colon | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Sigmoid Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 143.0 | 0 |
| Large Intestine, NOS | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Rectum | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Gallbladder | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1999

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 8.2 | 3.9 | 18.6 | 13 | 16.0 | 9.5 | 25.8 | 20 |
| Retroperitoneum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.6 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 8.9 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 89.7 | 73.0 | 111.2 | 126 | 83.6 | 68.2 | 102.0 | 113 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.6 | 0 |
| Larynx | 5.8 | 2.5 | 15.6 | 9 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 82.8 | 66.6 | 103.8 | 114 | 81.0 | 65.8 | 99.2 | 109 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.6 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 22.4 | 13.7 | 36.8 | 34 | 11.6 | 6.9 | 19.2 | 22 |
| Melanoma of the Skin | 19.3 | 11.2 | 33.3 | 29 | 9.3 | 5.4 | 16.1 | 19 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 144.8 | 126.1 | 166.2 | 252 |
| Female Genital System | 0.0 | 0.0 | 8.9 | 0 | 51.2 | 40.1 | 65.0 | 85 |
| Cervix Uteri | 0.0 | 0.0 | 8.9 | 0 | 4.7 | 2.0 | 10.7 | 9 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 8.9 | 0 | 21.8 | 15.1 | 31.3 | 38 |
| Corpus Uteri | 0.0 | 0.0 | 8.9 | 0 | 21.8 | 15.1 | 31.3 | 38 |
| Uterus, NOS | 0.0 | 0.0 | 8.9 | 0 | 0.0 | 0.0 | 4.6 | 0 |
| Ovary | 0.0 | 0.0 | 8.9 | 0 | 17.5 | 11.1 | 26.9 | 27 |
| Vagina | 0.0 | 0.0 | 8.9 | 0 | 0.0 | 0.0 | 4.6 | 0 |
| Vulva | 0.0 | 0.0 | 8.9 | 0 | 4.6 | 1.7 | 11.1 | 7 |
| Other Female Genital Organs | 0.0 | 0.0 | 8.9 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 189.7 | 164.3 | 219.9 | 261 | 0.0 | 0.0 | 4.6 | 0 |
| Prostate | 180.3 | 155.1 | 210.2 | 236 | 0.0 | 0.0 | 4.6 | 0 |
| Testis | 9.5 | 6.1 | 18.7 | 25 | 0.0 | 0.0 | 4.6 | 0 |
| Penis | 0.0 | 0.0 | 8.9 | 0 | 0.0 | 0.0 | 4.6 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 8.9 | 0 | 0.0 | 0.0 | 4.6 | 0 |
| Urinary System | 61.4 | 46.7 | 81.3 | 83 | 15.3 | 9.2 | 24.5 | 22 |
| Urinary Bladder | 40.2 | 27.9 | 58.0 | 47 | 9.2 | 4.4 | 17.4 | 11 |
| Kidney and Renal Pelvis | 20.4 | 13.2 | 33.0 | 35 | 6.1 | 2.9 | 12.6 | 11 |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.6 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 8.9 | 0 | 0.0 | 0.0 | 4.6 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1999

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Larynx | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Bones and Joints | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Melanoma of the Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Breast | 0.0 | 0.0 | 572.1 | 0 | 157.6 | 61.3 | 385.0 | 9 |
| Female Genital System | 0.0 | 0.0 | 572.1 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 572.1 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Corpus Uteri | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Uterus, NOS | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Ovary | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Vagina | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Vulva | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Male Genital System | 318.7 | 170.1 | 882.2 | 15 | 0.0 | 0.0 | 178.8 | 0 |
| Prostate | 318.7 | 170.1 | 882.2 | 15 | 0.0 | 0.0 | 178.8 | 0 |
| Testis | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Penis | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1999

| | American Indian/Alaska Native | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.6 | 0 |
| Pancreas | 31.4 | 9.4 | 84.8 | 6 | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.6 | 0 |
| Respiratory System | 101.5 | 63.6 | 164.2 | 25 | 62.7 | 36.9 | 102.5 | 19 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 39.5 | 0 | ^ | ^ | ^ | ^ |
| Larynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 94.1 | 57.5 | 155.7 | 23 | 51.3 | 28.0 | 89.0 | 15 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 39.5 | 0 | ^ | ^ | ^ | ^ |
| Bones and Joints | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 39.5 | 0 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 39.5 | 0 | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | 0.0 | 0.0 | 39.5 | 0 | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Breast | 0.0 | 0.0 | 39.5 | 0 | 87.4 | 58.8 | 128.6 | 32 |
| Female Genital System | 0.0 | 0.0 | 39.5 | 0 | 38.2 | 19.7 | 70.6 | 14 |
| Cervix Uteri | 0.0 | 0.0 | 39.5 | 0 | 18.5 | 6.7 | 45.1 | 7 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 39.5 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 39.5 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Ovary | 0.0 | 0.0 | 39.5 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 39.5 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Male Genital System | 81.8 | 48.7 | 140.0 | 21 | 0.0 | 0.0 | 19.6 | 0 |
| Prostate | 79.4 | 46.7 | 137.4 | 20 | 0.0 | 0.0 | 19.6 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.6 | 0 |
| Penis | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Urinary System | 77.9 | 41.1 | 142.3 | 17 | 32.5 | 14.4 | 65.6 | 9 |
| Urinary Bladder | 32.1 | 12.5 | 80.4 | 8 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 45.9 | 17.7 | 104.3 | 9 | 18.5 | 6.7 | 45.3 | 6 |
| Ureter | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1999

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Respiratory System | ^ | ^ | ^ | ^ | 49.8 | 17.7 | 193.5 | 6 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Larynx | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Bones and Joints | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Breast | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Female Genital System | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Ovary | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Vagina | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Vulva | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Male Genital System | 110.2 | 42.3 | 285.3 | 7 | 0.0 | 0.0 | 143.0 | 0 |
| Prostate | 110.2 | 42.3 | 285.3 | 7 | 0.0 | 0.0 | 143.0 | 0 |
| Testis | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Penis | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Urinary System | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Ureter | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1999

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.6 | 0 |
| Brain and Other Nervous System | 6.2 | 3.0 | 15.7 | 13 | 8.9 | 4.8 | 16.2 | 15 |
| Brain | 6.2 | 3.0 | 15.7 | 13 | 8.9 | 4.8 | 16.2 | 15 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 8.9 | 0 | 0.0 | 0.0 | 4.6 | 0 |
| Endocrine System | 8.3 | 4.7 | 17.9 | 19 | 8.7 | 5.2 | 15.1 | 20 |
| Thyroid | 8.0 | 4.4 | 17.6 | 18 | 8.7 | 5.2 | 15.1 | 20 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.6 | 0 |
| Lymphoma | 26.8 | 18.2 | 40.6 | 47 | 24.1 | 17.1 | 33.9 | 44 |
| Hodgkin Lymphoma | 2.9 | 1.2 | 11.6 | 8 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 2.9 | 1.2 | 11.6 | 8 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 8.9 | 0 | 0.0 | 0.0 | 4.6 | 0 |
| Non-Hodgkin Lymphoma | 23.9 | 15.6 | 37.6 | 39 | 21.9 | 15.2 | 31.5 | 39 |
| NHL - Nodal | 18.3 | 11.7 | 30.3 | 32 | 17.7 | 11.7 | 26.7 | 31 |
| NHL - Extranodal | 5.6 | 1.6 | 16.7 | 7 | 4.2 | 1.7 | 9.9 | 8 |
| Myeloma | 8.6 | 3.6 | 20.2 | 10 | ^ | ^ | ^ | ^ |
| Leukemia | 12.1 | 6.7 | 23.4 | 20 | 14.9 | 8.7 | 24.4 | 21 |
| Lymphocytic Leukemia | 8.6 | 4.3 | 19.0 | 15 | 6.2 | 2.5 | 13.5 | 8 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 6.6 | 3.3 | 16.2 | 13 | 5.4 | 1.9 | 12.6 | 6 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 8.9 | 0 | 0.0 | 0.0 | 4.6 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | 4.9 | 1.8 | 11.5 | 8 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | 3.3 | 1.0 | 9.1 | 6 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 8.9 | 0 | 0.0 | 0.0 | 4.6 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 8.9 | 0 | 0.0 | 0.0 | 4.6 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 8.9 | 0 | ^ | ^ | ^ | ^ |
| Mesothelioma | 8.0 | 2.9 | 19.9 | 8 | 0.0 | 0.0 | 4.6 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.6 | 0 |
| Miscellaneous | 5.0 | 2.1 | 14.5 | 8 | 9.3 | 4.5 | 17.7 | 11 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1999

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 572.1 | 0 | ^ | ^ | ^ | ^ |
| Brain | 0.0 | 0.0 | 572.1 | 0 | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Endocrine System | 0.0 | 0.0 | 572.1 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 572.1 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| NHL - Nodal | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| NHL - Extranodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Myeloma | 0.0 | 0.0 | 572.1 | 0 | ^ | ^ | ^ | ^ |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Chronic Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 572.1 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 572.1 | 0 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Leukemia | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Mesothelioma | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 572.1 | 0 | 0.0 | 0.0 | 178.8 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 178.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1999

| | American Indian/Alaska Native | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.6 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | 15.6 | 6.2 | 39.4 | 7 |
| Thyroid | ^ | ^ | ^ | ^ | 15.6 | 6.2 | 39.4 | 7 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Lymphoma | 12.6 | 4.5 | 51.9 | 6 | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Non-Hodgkin Lymphoma | 12.6 | 4.5 | 51.9 | 6 | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | 0.0 | 0.0 | 39.5 | 0 | ^ | ^ | ^ | ^ |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 39.5 | 0 | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 39.5 | 0 | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 39.5 | 0 | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.6 | 0 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.6 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.6 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Other Leukemia | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Mesothelioma | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 39.5 | 0 | 0.0 | 0.0 | 19.6 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

1999

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Endocrine System | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Lymphoma | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Non-Hodgkin Lymphoma | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| NHL - Nodal | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| NHL - Extranodal | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Myeloma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Leukemia | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 146.7 | 0 | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Other Leukemia | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Mesothelioma | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |
| Miscellaneous | 0.0 | 0.0 | 146.7 | 0 | 0.0 | 0.0 | 143.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2000

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 590.6 | 545.3 | 640.2 | 850 | 456.2 | 421.5 | 493.5 | 733 |
| Oral Cavity and Pharynx | 17.9 | 11.2 | 29.8 | 29 | 6.3 | 2.6 | 13.2 | 8 |
| Lip | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tongue | 6.4 | 2.6 | 16.3 | 11 | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.2 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.2 | 0 |
| Tonsil | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Oropharynx | 0.0 | 0.0 | 8.4 | 0 | ^ | ^ | ^ | ^ |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.2 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 8.4 | 0 | 0.0 | 0.0 | 4.2 | 0 |
| Digestive System | 115.4 | 95.9 | 139.5 | 163 | 78.7 | 63.6 | 96.6 | 104 |
| Esophagus | 12.3 | 6.7 | 23.5 | 18 | ^ | ^ | ^ | ^ |
| Stomach | 10.1 | 4.9 | 21.0 | 13 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 66.9 | 52.2 | 86.4 | 95 | 56.5 | 43.6 | 72.3 | 72 |
| Colon excluding Rectum | 49.0 | 35.9 | 67.1 | 62 | 41.2 | 30.2 | 55.2 | 51 |
| Cecum | 14.6 | 7.6 | 27.5 | 15 | 10.8 | 5.6 | 19.3 | 13 |
| Appendix | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ascending Colon | 7.8 | 3.4 | 18.0 | 10 | 8.6 | 4.0 | 16.5 | 10 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 8.4 | 0 | ^ | ^ | ^ | ^ |
| Descending Colon | 6.4 | 2.4 | 16.5 | 8 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 14.6 | 7.9 | 27.0 | 19 | 11.8 | 6.4 | 20.4 | 15 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 17.9 | 11.8 | 29.1 | 33 | 15.3 | 9.2 | 24.4 | 21 |
| Rectosigmoid Junction | 4.0 | 1.6 | 12.6 | 8 | ^ | ^ | ^ | ^ |
| Rectum | 14.0 | 8.5 | 24.7 | 25 | 12.0 | 6.7 | 20.3 | 17 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | 11.6 | 6.4 | 22.2 | 18 | ^ | ^ | ^ | ^ |
| Liver | 10.3 | 5.4 | 20.7 | 16 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gallbladder | 0.0 | 0.0 | 8.4 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2000

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 718.4 | 472.2 | 1,212.1 | 35 | 238.6 | 125.4 | 457.3 | 16 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Lip | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Tongue | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Salivary Gland | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Nasopharynx | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Tonsil | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Oropharynx | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Hypopharynx | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Digestive System | 188.9 | 76.4 | 607.3 | 10 | ^ | ^ | ^ | ^ |
| Esophagus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 156.7 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 156.7 | 0 |
| Small Intestine | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 156.7 | 0 |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 156.7 | 0 |
| Cecum | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Appendix | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Ascending Colon | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Transverse Colon | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Descending Colon | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Sigmoid Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 156.7 | 0 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 156.7 | 0 |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Rectosigmoid Junction | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Rectum | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Liver and Intrahepatic Bile Duct | 0.0 | 0.0 | 408.6 | 0 | ^ | ^ | ^ | ^ |
| Liver | 0.0 | 0.0 | 408.6 | 0 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Gallbladder | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2000

| | American Indian/Alaska Native | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 490.2 | 400.4 | 600.8 | 130 | 499.8 | 423.0 | 588.9 | 165 |
| Oral Cavity and Pharynx | 25.5 | 9.9 | 67.6 | 7 | ^ | ^ | ^ | ^ |
| Lip | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.2 | 0 |
| Tongue | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.2 | 0 |
| Salivary Gland | 0.0 | 0.0 | 36.2 | 0 | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.2 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.2 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Oropharynx | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.2 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Digestive System | 169.2 | 118.6 | 242.0 | 45 | 170.1 | 124.3 | 229.2 | 49 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | 41.0 | 16.8 | 92.0 | 9 | 19.1 | 6.9 | 45.9 | 6 |
| Small Intestine | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Colon and Rectum | 94.7 | 57.7 | 155.2 | 25 | 100.5 | 65.2 | 149.9 | 27 |
| Colon excluding Rectum | 54.4 | 27.5 | 106.1 | 14 | 72.1 | 42.7 | 115.8 | 19 |
| Cecum | ^ | ^ | ^ | ^ | 20.7 | 7.3 | 49.3 | 6 |
| Appendix | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Ascending Colon | 0.0 | 0.0 | 36.2 | 0 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.2 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | 24.6 | 8.8 | 56.2 | 6 |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Sigmoid Colon | 28.2 | 10.2 | 73.3 | 7 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.2 | 0 |
| Rectum and Rectosigmoid Junction | 40.3 | 17.8 | 88.6 | 11 | 28.4 | 11.6 | 60.4 | 8 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 37.0 | 15.5 | 84.8 | 10 | 23.2 | 8.8 | 52.6 | 7 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.2 | 0 |
| Liver | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.2 | 0 |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.2 | 0 |
| Gallbladder | 0.0 | 0.0 | 36.2 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2000

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 400.1 | 249.1 | 632.3 | 28 | 360.9 | 255.5 | 526.3 | 47 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Lip | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Tongue | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Salivary Gland | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Nasopharynx | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Tonsil | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Oropharynx | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Hypopharynx | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Digestive System | 166.6 | 79.0 | 340.0 | 12 | 116.2 | 52.8 | 253.0 | 12 |
| Esophagus | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Stomach | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | 53.0 | 19.3 | 164.6 | 6 |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Ascending Colon | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Transverse Colon | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Descending Colon | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Sigmoid Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 104.0 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Gallbladder | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2000

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 7.0 | 2.9 | 17.1 | 10 | 7.9 | 3.8 | 15.0 | 11 |
| Retroperitoneum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.2 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 8.4 | 0 | 0.0 | 0.0 | 4.2 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.2 | 0 |
| Respiratory System | 91.4 | 73.7 | 113.7 | 121 | 54.7 | 43.0 | 69.2 | 81 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.2 | 0 |
| Larynx | 4.2 | 1.7 | 13.0 | 8 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 85.6 | 68.4 | 107.6 | 111 | 52.9 | 41.3 | 67.2 | 78 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | 4.3 | 1.4 | 13.4 | 8 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 13.6 | 7.9 | 24.7 | 28 | 12.2 | 7.3 | 20.1 | 22 |
| Melanoma of the Skin | 13.6 | 7.9 | 24.7 | 28 | 11.2 | 6.6 | 18.6 | 21 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 8.4 | 0 | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 154.7 | 135.7 | 176.1 | 273 |
| Female Genital System | 0.0 | 0.0 | 8.4 | 0 | 50.4 | 40.0 | 63.3 | 92 |
| Cervix Uteri | 0.0 | 0.0 | 8.4 | 0 | 6.4 | 3.6 | 12.0 | 15 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 8.4 | 0 | 23.6 | 16.8 | 33.0 | 44 |
| Corpus Uteri | 0.0 | 0.0 | 8.4 | 0 | 23.2 | 16.5 | 32.7 | 43 |
| Uterus, NOS | 0.0 | 0.0 | 8.4 | 0 | ^ | ^ | ^ | ^ |
| Ovary | 0.0 | 0.0 | 8.4 | 0 | 15.6 | 9.9 | 24.2 | 26 |
| Vagina | 0.0 | 0.0 | 8.4 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 8.4 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 8.4 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 200.3 | 174.3 | 230.7 | 275 | 0.0 | 0.0 | 4.2 | 0 |
| Prostate | 193.7 | 168.0 | 223.9 | 260 | 0.0 | 0.0 | 4.2 | 0 |
| Testis | 4.5 | 2.3 | 12.7 | 12 | 0.0 | 0.0 | 4.2 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.2 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 8.4 | 0 | 0.0 | 0.0 | 4.2 | 0 |
| Urinary System | 56.6 | 42.0 | 76.4 | 75 | 24.1 | 15.9 | 35.3 | 30 |
| Urinary Bladder | 41.8 | 28.9 | 60.0 | 51 | 14.0 | 7.8 | 23.3 | 16 |
| Kidney and Renal Pelvis | 13.8 | 7.8 | 25.2 | 23 | 10.1 | 5.3 | 18.1 | 14 |
| Ureter | 0.0 | 0.0 | 8.4 | 0 | 0.0 | 0.0 | 4.2 | 0 |
| Other Urinary Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2000

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 156.7 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 156.7 | 0 |
| Respiratory System | 192.6 | 70.9 | 621.6 | 7 | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 156.7 | 0 |
| Lung and Bronchus | 150.1 | 50.0 | 567.7 | 6 | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Bones and Joints | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Breast | 0.0 | 0.0 | 408.6 | 0 | 119.5 | 40.7 | 317.1 | 7 |
| Female Genital System | 0.0 | 0.0 | 408.6 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 408.6 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 408.6 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 408.6 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Ovary | 0.0 | 0.0 | 408.6 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 408.6 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Male Genital System | 324.1 | 172.9 | 755.4 | 16 | 0.0 | 0.0 | 156.7 | 0 |
| Prostate | 324.1 | 172.9 | 755.4 | 16 | 0.0 | 0.0 | 156.7 | 0 |
| Testis | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Penis | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Urinary System | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Urinary Bladder | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Kidney and Renal Pelvis | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Ureter | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2000

| | American Indian/Alaska Native | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | ^ | ^ | ^ | ^ | 32.0 | 15.4 | 62.5 | 11 |
| Retroperitoneum | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Respiratory System | 129.9 | 84.3 | 198.9 | 31 | 77.7 | 47.3 | 122.4 | 21 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.2 | 0 |
| Lung and Bronchus | 118.8 | 76.0 | 184.9 | 29 | 72.5 | 43.5 | 115.6 | 20 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.2 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.2 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Breast | 0.0 | 0.0 | 36.2 | 0 | 153.8 | 116.4 | 202.9 | 60 |
| Female Genital System | 0.0 | 0.0 | 36.2 | 0 | 22.7 | 9.5 | 50.1 | 8 |
| Cervix Uteri | 0.0 | 0.0 | 36.2 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 36.2 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 36.2 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Ovary | 0.0 | 0.0 | 36.2 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Vulva | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Male Genital System | 68.2 | 36.5 | 124.8 | 17 | 0.0 | 0.0 | 19.2 | 0 |
| Prostate | 64.1 | 33.0 | 120.5 | 15 | 0.0 | 0.0 | 19.2 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.2 | 0 |
| Penis | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Urinary System | 45.1 | 21.5 | 93.9 | 13 | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 24.6 | 10.3 | 64.9 | 8 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2000

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Larynx | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Bones and Joints | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 104.0 | 0 |
| Breast | 0.0 | 0.0 | 121.7 | 0 | 80.3 | 42.1 | 190.3 | 13 |
| Female Genital System | 0.0 | 0.0 | 121.7 | 0 | 45.8 | 17.8 | 152.2 | 7 |
| Cervix Uteri | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Ovary | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Vulva | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Male Genital System | 94.5 | 31.3 | 250.8 | 6 | 0.0 | 0.0 | 104.0 | 0 |
| Prostate | 94.5 | 31.3 | 250.8 | 6 | 0.0 | 0.0 | 104.0 | 0 |
| Testis | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Penis | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2000

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 8.4 | 0 | ^ | ^ | ^ | ^ |
| Brain and Other Nervous System | 7.1 | 3.9 | 15.8 | 16 | 7.6 | 4.0 | 14.2 | 14 |
| Brain | 6.4 | 3.5 | 15.0 | 15 | 7.6 | 4.0 | 14.2 | 14 |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.2 | 0 |
| Endocrine System | 6.8 | 3.7 | 15.7 | 15 | 8.3 | 5.1 | 14.2 | 20 |
| Thyroid | 6.0 | 3.1 | 14.8 | 13 | 8.3 | 5.1 | 14.2 | 20 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 4.2 | 0 |
| Lymphoma | 26.6 | 18.7 | 39.3 | 48 | 21.5 | 14.2 | 31.7 | 32 |
| Hodgkin Lymphoma | 3.9 | 1.4 | 12.6 | 7 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 3.9 | 1.4 | 12.6 | 7 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 8.4 | 0 | 0.0 | 0.0 | 4.2 | 0 |
| Non-Hodgkin Lymphoma | 22.7 | 15.4 | 35.0 | 41 | 20.6 | 13.5 | 30.7 | 30 |
| NHL - Nodal | 17.1 | 10.7 | 28.8 | 29 | 13.0 | 7.5 | 21.4 | 19 |
| NHL - Extranodal | 5.6 | 2.7 | 14.4 | 12 | 7.7 | 3.5 | 15.0 | 11 |
| Myeloma | 5.6 | 2.4 | 14.6 | 10 | 5.3 | 1.9 | 12.0 | 6 |
| Leukemia | 25.1 | 16.2 | 39.1 | 35 | 15.6 | 9.7 | 24.4 | 25 |
| Lymphocytic Leukemia | 10.2 | 5.6 | 20.3 | 18 | 5.8 | 2.5 | 12.2 | 9 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 9.1 | 4.6 | 19.1 | 15 | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloid and Monocytic Leukemia | 14.9 | 7.8 | 27.6 | 17 | 8.5 | 4.2 | 15.9 | 13 |
| Acute Myeloid Leukemia | 10.3 | 4.6 | 21.9 | 12 | 6.3 | 2.6 | 13.3 | 9 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 8.4 | 0 | ^ | ^ | ^ | ^ |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Leukemia | 0.0 | 0.0 | 8.4 | 0 | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | 0.0 | 0.0 | 8.4 | 0 | ^ | ^ | ^ | ^ |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 8.4 | 0 | ^ | ^ | ^ | ^ |
| Mesothelioma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kaposi Sarcoma | 0.0 | 0.0 | 8.4 | 0 | 0.0 | 0.0 | 4.2 | 0 |
| Miscellaneous | 16.2 | 9.0 | 28.9 | 21 | 10.6 | 5.6 | 18.8 | 14 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2000

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Brain | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Endocrine System | 0.0 | 0.0 | 408.6 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 408.6 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 156.7 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 156.7 | 0 |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 156.7 | 0 |
| NHL - Extranodal | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Myeloma | 0.0 | 0.0 | 408.6 | 0 | ^ | ^ | ^ | ^ |
| Leukemia | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Lymphocytic Leukemia | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Other Leukemia | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Mesothelioma | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 408.6 | 0 | 0.0 | 0.0 | 156.7 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2000

| | American Indian/Alaska Native | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.2 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Brain | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Endocrine System | 0.0 | 0.0 | 36.2 | 0 | 14.7 | 5.8 | 37.6 | 7 |
| Thyroid | 0.0 | 0.0 | 36.2 | 0 | 14.7 | 5.8 | 37.6 | 7 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Lymphoma | 17.5 | 5.4 | 57.2 | 6 | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 36.2 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 0.0 | 0.0 | 36.2 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Non-Hodgkin Lymphoma | 17.5 | 5.4 | 57.2 | 6 | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Leukemia | 0.0 | 0.0 | 36.2 | 0 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 36.2 | 0 | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 36.2 | 0 | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 36.2 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 36.2 | 0 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Other Leukemia | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Mesothelioma | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 36.2 | 0 | 0.0 | 0.0 | 19.2 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2000

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Brain | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Endocrine System | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 104.0 | 0 |
| NHL - Extranodal | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 104.0 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 104.0 | 0 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 104.0 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 121.7 | 0 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Other Leukemia | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Mesothelioma | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 121.7 | 0 | 0.0 | 0.0 | 104.0 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 104.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2001

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 601.2 | 556.5 | 650.2 | 918 | 435.2 | 402.5 | 470.4 | 747 |
| Oral Cavity and Pharynx | 16.7 | 10.8 | 27.7 | 32 | 4.6 | 1.9 | 10.2 | 8 |
| Lip | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.9 | 0 |
| Tongue | 4.9 | 1.8 | 14.2 | 8 | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.9 | 0 |
| Tonsil | 4.5 | 2.0 | 13.0 | 10 | ^ | ^ | ^ | ^ |
| Oropharynx | 0.0 | 0.0 | 8.3 | 0 | 0.0 | 0.0 | 3.9 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.9 | 0 |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Digestive System | 100.2 | 82.1 | 122.9 | 150 | 79.5 | 65.0 | 96.7 | 115 |
| Esophagus | 7.1 | 3.2 | 16.9 | 12 | 5.4 | 2.1 | 11.9 | 7 |
| Stomach | 7.9 | 4.1 | 17.3 | 16 | 5.4 | 2.1 | 11.8 | 7 |
| Small Intestine | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.9 | 0 |
| Colon and Rectum | 67.2 | 51.9 | 87.3 | 89 | 41.4 | 31.1 | 54.5 | 59 |
| Colon excluding Rectum | 51.8 | 38.1 | 70.6 | 64 | 28.1 | 19.6 | 39.4 | 38 |
| Cecum | 7.5 | 3.6 | 17.1 | 11 | 10.4 | 5.6 | 18.1 | 15 |
| Appendix | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.9 | 0 |
| Ascending Colon | 16.1 | 8.1 | 30.0 | 13 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | 13.8 | 7.6 | 25.6 | 19 | 5.7 | 2.4 | 12.1 | 8 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 15.4 | 9.4 | 26.6 | 25 | 13.3 | 8.0 | 21.5 | 21 |
| Rectosigmoid Junction | 4.0 | 1.3 | 13.2 | 6 | ^ | ^ | ^ | ^ |
| Rectum | 11.3 | 6.3 | 21.7 | 19 | 12.5 | 7.3 | 20.6 | 19 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 8.3 | 0 | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | 8.0 | 3.5 | 18.4 | 14 | 6.0 | 2.7 | 12.4 | 10 |
| Liver | 6.8 | 2.6 | 17.2 | 11 | 5.3 | 2.1 | 11.6 | 8 |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2001

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 683.2 | 445.5 | 1,137.1 | 35 | 482.6 | 283.2 | 798.5 | 26 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Lip | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Tongue | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Salivary Gland | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Nasopharynx | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Tonsil | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Oropharynx | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Hypopharynx | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Digestive System | 128.1 | 40.5 | 494.9 | 7 | 179.9 | 66.3 | 423.7 | 7 |
| Esophagus | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| Small Intestine | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | 162.9 | 54.6 | 404.3 | 6 |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 357.6 | 0 | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| Transverse Colon | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Descending Colon | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 357.6 | 0 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| Liver | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Gallbladder | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2001

| | American Indian/Alaska Native | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 544.8 | 449.6 | 663.8 | 145 | 495.1 | 421.6 | 580.3 | 175 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 23.7 | 9.8 | 52.0 | 8 |
| Lip | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Tongue | 0.0 | 0.0 | 43.7 | 0 | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 43.7 | 0 | ^ | ^ | ^ | ^ |
| Floor of Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Oropharynx | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Hypopharynx | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Digestive System | 199.0 | 145.0 | 278.1 | 53 | 172.2 | 128.2 | 228.7 | 55 |
| Esophagus | 29.9 | 10.0 | 82.7 | 6 | ^ | ^ | ^ | ^ |
| Stomach | 24.4 | 8.0 | 73.8 | 6 | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 43.7 | 0 | ^ | ^ | ^ | ^ |
| Colon and Rectum | 104.3 | 66.8 | 168.5 | 28 | 108.8 | 75.2 | 155.1 | 36 |
| Colon excluding Rectum | 49.4 | 25.5 | 102.6 | 13 | 89.1 | 58.4 | 132.8 | 28 |
| Cecum | ^ | ^ | ^ | ^ | 16.2 | 5.9 | 40.6 | 6 |
| Appendix | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 43.7 | 0 | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Sigmoid Colon | ^ | ^ | ^ | ^ | 39.6 | 19.8 | 73.5 | 12 |
| Large Intestine, NOS | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Rectum and Rectosigmoid Junction | 54.9 | 28.4 | 110.6 | 15 | 19.7 | 8.2 | 44.9 | 8 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 36.5 | 16.6 | 86.9 | 11 | 17.5 | 6.8 | 42.3 | 7 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | 19.2 | 6.3 | 65.5 | 6 | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.0 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2001

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 395.8 | 259.8 | 611.9 | 33 | 264.8 | 175.9 | 407.8 | 34 |
| Oral Cavity and Pharynx | 0.0 | 0.0 | 130.0 | 0 | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Tongue | 0.0 | 0.0 | 130.0 | 0 | ^ | ^ | ^ | ^ |
| Salivary Gland | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Nasopharynx | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Tonsil | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Oropharynx | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Hypopharynx | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Digestive System | 140.3 | 60.7 | 313.2 | 11 | 51.1 | 17.8 | 151.1 | 6 |
| Esophagus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 89.4 | 0 |
| Stomach | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 89.4 | 0 |
| Small Intestine | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 89.4 | 0 |
| Appendix | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Ascending Colon | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Transverse Colon | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Descending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 89.4 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 130.0 | 0 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Gallbladder | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2001

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 6.0 | 2.8 | 15.0 | 12 | 13.7 | 8.1 | 22.1 | 19 |
| Retroperitoneum | 0.0 | 0.0 | 8.3 | 0 | ^ | ^ | ^ | ^ |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 8.3 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.9 | 0 |
| Respiratory System | 88.9 | 71.6 | 110.7 | 127 | 63.9 | 51.2 | 79.2 | 97 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Larynx | 6.9 | 3.0 | 16.7 | 10 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 80.0 | 63.5 | 101.1 | 113 | 61.2 | 48.7 | 76.3 | 91 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.9 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 16.7 | 11.1 | 27.3 | 35 | 16.6 | 10.8 | 25.1 | 30 |
| Melanoma of the Skin | 16.4 | 10.8 | 27.0 | 34 | 15.8 | 10.1 | 24.1 | 28 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 139.4 | 121.9 | 159.3 | 261 |
| Female Genital System | 0.0 | 0.0 | 8.3 | 0 | 46.1 | 36.2 | 58.4 | 85 |
| Cervix Uteri | 0.0 | 0.0 | 8.3 | 0 | 9.2 | 5.4 | 15.7 | 19 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 8.3 | 0 | 25.1 | 17.9 | 34.8 | 46 |
| Corpus Uteri | 0.0 | 0.0 | 8.3 | 0 | 23.7 | 16.8 | 33.1 | 44 |
| Uterus, NOS | 0.0 | 0.0 | 8.3 | 0 | ^ | ^ | ^ | ^ |
| Ovary | 0.0 | 0.0 | 8.3 | 0 | 8.2 | 4.3 | 15.0 | 14 |
| Vagina | 0.0 | 0.0 | 8.3 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 8.3 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 8.3 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 204.2 | 179.0 | 233.7 | 311 | 0.0 | 0.0 | 3.9 | 0 |
| Prostate | 196.2 | 171.2 | 225.4 | 291 | 0.0 | 0.0 | 3.9 | 0 |
| Testis | 7.6 | 4.5 | 16.2 | 19 | 0.0 | 0.0 | 3.9 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.9 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 8.3 | 0 | 0.0 | 0.0 | 3.9 | 0 |
| Urinary System | 75.1 | 59.3 | 95.7 | 105 | 15.8 | 10.1 | 24.3 | 26 |
| Urinary Bladder | 51.2 | 38.4 | 68.8 | 71 | 8.0 | 4.1 | 14.7 | 13 |
| Kidney and Renal Pelvis | 22.4 | 14.0 | 36.2 | 32 | 6.5 | 3.1 | 12.8 | 11 |
| Ureter | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Urinary Organs | 0.0 | 0.0 | 8.3 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2001

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Pancreas | 0.0 | 0.0 | 357.6 | 0 | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Peritoneum, Omentum and Mesentery | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Larynx | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Bones and Joints | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Breast | 0.0 | 0.0 | 357.6 | 0 | 155.2 | 53.9 | 386.4 | 9 |
| Female Genital System | 0.0 | 0.0 | 357.6 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 357.6 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 357.6 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 357.6 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Ovary | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Vagina | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Vulva | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Male Genital System | 300.0 | 147.7 | 698.7 | 14 | 0.0 | 0.0 | 168.8 | 0 |
| Prostate | 300.0 | 147.7 | 698.7 | 14 | 0.0 | 0.0 | 168.8 | 0 |
| Testis | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Penis | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| Ureter | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2001

| | American Indian/Alaska Native | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 43.7 | 0 | ^ | ^ | ^ | ^ |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Respiratory System | 147.5 | 98.8 | 223.1 | 35 | 69.8 | 43.1 | 109.8 | 22 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.0 | 0 |
| Larynx | 0.0 | 0.0 | 43.7 | 0 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 144.8 | 96.3 | 220.1 | 34 | 67.0 | 40.8 | 106.5 | 21 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Bones and Joints | 0.0 | 0.0 | 43.7 | 0 | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.0 | 0 |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.0 | 0 |
| Breast | 0.0 | 0.0 | 43.7 | 0 | 132.3 | 97.7 | 178.6 | 52 |
| Female Genital System | 0.0 | 0.0 | 43.7 | 0 | 37.4 | 20.8 | 66.8 | 15 |
| Cervix Uteri | 0.0 | 0.0 | 43.7 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 43.7 | 0 | 19.9 | 8.6 | 45.0 | 8 |
| Corpus Uteri | 0.0 | 0.0 | 43.7 | 0 | 19.9 | 8.6 | 45.0 | 8 |
| Uterus, NOS | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Ovary | 0.0 | 0.0 | 43.7 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Vulva | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 43.7 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 96.9 | 57.4 | 164.6 | 26 | 0.0 | 0.0 | 19.0 | 0 |
| Prostate | 89.5 | 50.9 | 156.9 | 22 | 0.0 | 0.0 | 19.0 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.0 | 0 |
| Penis | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Urinary System | 41.8 | 21.3 | 91.9 | 12 | ^ | ^ | ^ | ^ |
| Urinary Bladder | 21.5 | 7.7 | 68.1 | 6 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 20.3 | 7.3 | 66.2 | 6 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2001

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 89.4 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Larynx | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 89.4 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 89.4 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Breast | 0.0 | 0.0 | 130.0 | 0 | 78.2 | 39.4 | 178.2 | 12 |
| Female Genital System | 0.0 | 0.0 | 130.0 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 130.0 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 130.0 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 130.0 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Ovary | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Vagina | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Vulva | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Male Genital System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 89.4 | 0 |
| Prostate | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 89.4 | 0 |
| Testis | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Penis | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 130.0 | 0 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Other Urinary Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 89.4 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2001

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.9 | 0 |
| Brain and Other Nervous System | 11.4 | 6.7 | 21.4 | 22 | 5.3 | 2.4 | 11.0 | 10 |
| Brain | 11.4 | 6.7 | 21.4 | 22 | 4.1 | 1.7 | 9.3 | 8 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 8.3 | 0 | ^ | ^ | ^ | ^ |
| Endocrine System | 5.1 | 2.6 | 13.4 | 13 | 15.9 | 10.8 | 23.5 | 34 |
| Thyroid | 3.6 | 1.7 | 11.8 | 10 | 12.7 | 8.3 | 19.5 | 28 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 3.2 | 1.0 | 8.4 | 6 |
| Lymphoma | 28.5 | 19.8 | 42.1 | 49 | 18.5 | 12.3 | 27.3 | 32 |
| Hodgkin Lymphoma | 3.2 | 1.2 | 11.5 | 7 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 3.2 | 1.2 | 11.5 | 7 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 8.3 | 0 | 0.0 | 0.0 | 3.9 | 0 |
| Non-Hodgkin Lymphoma | 25.4 | 17.0 | 38.7 | 42 | 17.2 | 11.2 | 25.9 | 29 |
| NHL - Nodal | 15.4 | 9.7 | 26.3 | 29 | 13.5 | 8.3 | 21.4 | 23 |
| NHL - Extranodal | 9.9 | 4.5 | 21.2 | 13 | 3.7 | 1.3 | 9.1 | 6 |
| Myeloma | 5.8 | 2.3 | 15.3 | 8 | ^ | ^ | ^ | ^ |
| Leukemia | 14.7 | 8.0 | 27.1 | 19 | 12.6 | 7.5 | 20.4 | 21 |
| Lymphocytic Leukemia | 7.0 | 3.1 | 16.7 | 11 | 5.1 | 2.1 | 11.1 | 8 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 5.0 | 1.6 | 14.7 | 6 | 4.3 | 1.5 | 10.2 | 6 |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.9 | 0 |
| Myeloid and Monocytic Leukemia | 6.9 | 2.3 | 18.0 | 7 | 6.5 | 3.2 | 12.6 | 12 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | 3.6 | 1.2 | 9.0 | 6 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 8.3 | 0 | ^ | ^ | ^ | ^ |
| Other Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.9 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 8.3 | 0 | ^ | ^ | ^ | ^ |
| Mesothelioma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.9 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 8.3 | 0 | ^ | ^ | ^ | ^ |
| Miscellaneous | 22.2 | 13.9 | 35.7 | 31 | 11.1 | 6.2 | 18.8 | 17 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2001

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Brain | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Endocrine System | 0.0 | 0.0 | 357.6 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 357.6 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| NHL - Extranodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |
| Myeloma | 0.0 | 0.0 | 357.6 | 0 | ^ | ^ | ^ | ^ |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 357.6 | 0 | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Other Leukemia | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Mesothelioma | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 357.6 | 0 | 0.0 | 0.0 | 168.8 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 168.8 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2001

| | American Indian/Alaska Native | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 43.7 | 0 | ^ | ^ | ^ | ^ |
| Brain | 0.0 | 0.0 | 43.7 | 0 | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 43.7 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.0 | 0 |
| Lymphoma | 20.2 | 7.9 | 65.1 | 7 | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.0 | 0 |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.0 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Non-Hodgkin Lymphoma | 17.4 | 6.2 | 62.1 | 6 | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | 0.0 | 0.0 | 43.7 | 0 | ^ | ^ | ^ | ^ |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.0 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 19.0 | 0 |
| Mesothelioma | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 43.7 | 0 | 0.0 | 0.0 | 19.0 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2001

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Brain | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 89.4 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 89.4 | 0 |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 89.4 | 0 |
| NHL - Extranodal | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 89.4 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Other Leukemia | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Mesothelioma | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 130.0 | 0 | 0.0 | 0.0 | 89.4 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 89.4 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2002

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 586.1 | 543.6 | 632.5 | 957 | 382.4 | 352.7 | 414.3 | 698 |
| Oral Cavity and Pharynx | 13.1 | 8.3 | 22.7 | 29 | 5.7 | 2.5 | 11.6 | 9 |
| Lip | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Tongue | 3.3 | 1.2 | 11.1 | 7 | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Oropharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Digestive System | 99.3 | 81.9 | 120.9 | 158 | 65.0 | 52.2 | 80.1 | 101 |
| Esophagus | 10.5 | 5.2 | 21.2 | 15 | ^ | ^ | ^ | ^ |
| Stomach | 7.3 | 2.9 | 17.2 | 10 | ^ | ^ | ^ | ^ |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 58.9 | 45.7 | 76.5 | 90 | 39.1 | 29.2 | 51.5 | 58 |
| Colon excluding Rectum | 38.8 | 27.7 | 54.4 | 54 | 30.0 | 21.4 | 41.1 | 44 |
| Cecum | 9.3 | 3.6 | 20.9 | 8 | 7.4 | 3.5 | 13.9 | 11 |
| Appendix | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Ascending Colon | 5.5 | 2.3 | 14.1 | 9 | 5.5 | 2.3 | 11.6 | 8 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | 5.6 | 2.2 | 11.9 | 7 |
| Transverse Colon | 3.6 | 1.1 | 12.0 | 6 | ^ | ^ | ^ | ^ |
| Splenic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Sigmoid Colon | 12.6 | 6.7 | 23.6 | 18 | 8.1 | 4.3 | 14.6 | 14 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 20.1 | 13.4 | 31.4 | 36 | 9.1 | 4.8 | 16.1 | 14 |
| Rectosigmoid Junction | 4.3 | 1.6 | 12.6 | 8 | ^ | ^ | ^ | ^ |
| Rectum | 15.9 | 10.0 | 26.4 | 28 | 7.8 | 3.7 | 14.6 | 11 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | 2.9 | 1.0 | 7.4 | 6 |
| Liver and Intrahepatic Bile Duct | 5.8 | 2.8 | 14.1 | 13 | ^ | ^ | ^ | ^ |
| Liver | 5.1 | 2.3 | 13.3 | 11 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Gallbladder | 0.0 | 0.0 | 7.7 | 0 | 0.0 | 0.0 | 3.5 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2002

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 483.6 | 312.7 | 829.3 | 32 | 399.0 | 243.2 | 650.5 | 26 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Tongue | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Salivary Gland | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |
| Floor of Mouth | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 146.3 | 0 |
| Tonsil | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Oropharynx | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Hypopharynx | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Digestive System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Esophagus | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Stomach | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Small Intestine | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Appendix | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Ascending Colon | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Hepatic Flexure | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 146.3 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 146.3 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Descending Colon | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Large Intestine, NOS | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Rectosigmoid Junction | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Rectum | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 146.3 | 0 |
| Liver | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 146.3 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Gallbladder | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2002

| | American Indian/Alaska Native | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 520.0 | 429.5 | 634.4 | 153 | 507.5 | 430.9 | 595.8 | 171 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Tongue | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Salivary Gland | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 18.1 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Tonsil | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Oropharynx | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Hypopharynx | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Digestive System | 204.0 | 142.8 | 292.3 | 47 | 173.8 | 128.6 | 231.5 | 53 |
| Esophagus | 28.8 | 9.6 | 81.8 | 6 | ^ | ^ | ^ | ^ |
| Stomach | 57.8 | 26.0 | 121.9 | 12 | 32.5 | 15.0 | 63.9 | 10 |
| Small Intestine | 0.0 | 0.0 | 44.7 | 0 | ^ | ^ | ^ | ^ |
| Colon and Rectum | 64.1 | 35.5 | 121.8 | 17 | 109.6 | 74.3 | 157.7 | 33 |
| Colon excluding Rectum | 41.8 | 18.5 | 96.7 | 10 | 74.8 | 46.1 | 116.9 | 22 |
| Cecum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Appendix | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | 0.0 | 0.0 | 44.7 | 0 | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Descending Colon | 0.0 | 0.0 | 44.7 | 0 | ^ | ^ | ^ | ^ |
| Sigmoid Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 22.3 | 8.8 | 69.0 | 7 | 34.7 | 16.8 | 66.1 | 11 |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 18.1 | 0 |
| Rectum | ^ | ^ | ^ | ^ | 34.7 | 16.8 | 66.1 | 11 |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 44.7 | 0 | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2002

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 396.1 | 248.5 | 626.4 | 32 | 335.9 | 240.6 | 474.9 | 50 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 74.3 | 0 |
| Lip | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Tongue | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Salivary Gland | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 74.3 | 0 |
| Tonsil | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Oropharynx | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Hypopharynx | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Digestive System | 54.7 | 16.7 | 190.8 | 6 | 79.4 | 34.7 | 177.2 | 11 |
| Esophagus | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Stomach | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | 48.3 | 13.5 | 141.7 | 6 |
| Colon excluding Rectum | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Appendix | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Ascending Colon | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Transverse Colon | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Descending Colon | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Gallbladder | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2002

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 12.7 | 7.1 | 23.3 | 22 | 10.0 | 5.5 | 17.1 | 16 |
| Retroperitoneum | 0.0 | 0.0 | 7.7 | 0 | 0.0 | 0.0 | 3.5 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 7.7 | 0 | 0.0 | 0.0 | 3.5 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 90.3 | 73.4 | 111.5 | 130 | 55.1 | 43.8 | 68.9 | 89 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Larynx | 3.3 | 1.1 | 11.3 | 7 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 85.2 | 68.6 | 106.0 | 120 | 53.0 | 41.9 | 66.6 | 86 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | 3.9 | 1.3 | 12.2 | 7 | 2.8 | 1.0 | 7.2 | 6 |
| Skin excluding Basal and Squamous | 18.5 | 12.0 | 29.5 | 38 | 14.3 | 9.5 | 21.4 | 31 |
| Melanoma of the Skin | 17.2 | 11.0 | 28.1 | 35 | 13.9 | 9.2 | 20.9 | 30 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 125.3 | 109.2 | 143.6 | 245 |
| Female Genital System | 0.0 | 0.0 | 7.7 | 0 | 44.5 | 35.0 | 56.3 | 86 |
| Cervix Uteri | 0.0 | 0.0 | 7.7 | 0 | 8.5 | 4.9 | 14.5 | 19 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 7.7 | 0 | 18.1 | 12.1 | 26.6 | 32 |
| Corpus Uteri | 0.0 | 0.0 | 7.7 | 0 | 16.8 | 11.0 | 24.9 | 30 |
| Uterus, NOS | 0.0 | 0.0 | 7.7 | 0 | ^ | ^ | ^ | ^ |
| Ovary | 0.0 | 0.0 | 7.7 | 0 | 12.0 | 7.3 | 19.0 | 23 |
| Vagina | 0.0 | 0.0 | 7.7 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 7.7 | 0 | 4.7 | 2.0 | 10.1 | 9 |
| Other Female Genital Organs | 0.0 | 0.0 | 7.7 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 206.4 | 182.0 | 234.7 | 339 | 0.0 | 0.0 | 3.5 | 0 |
| Prostate | 197.6 | 173.5 | 225.6 | 317 | 0.0 | 0.0 | 3.5 | 0 |
| Testis | 8.5 | 5.2 | 16.8 | 21 | 0.0 | 0.0 | 3.5 | 0 |
| Penis | 0.0 | 0.0 | 7.7 | 0 | 0.0 | 0.0 | 3.5 | 0 |
| Other Male Genital Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Urinary System | 71.4 | 56.8 | 90.1 | 110 | 12.7 | 7.7 | 20.3 | 22 |
| Urinary Bladder | 51.7 | 39.2 | 68.8 | 73 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 19.6 | 13.0 | 30.9 | 37 | 8.9 | 4.8 | 15.5 | 16 |
| Ureter | 0.0 | 0.0 | 7.7 | 0 | ^ | ^ | ^ | ^ |
| Other Urinary Organs | 0.0 | 0.0 | 7.7 | 0 | 0.0 | 0.0 | 3.5 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2002

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 146.3 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Larynx | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Bones and Joints | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Breast | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |
| Female Genital System | 0.0 | 0.0 | 286.0 | 0 | 105.2 | 37.6 | 281.8 | 7 |
| Cervix Uteri | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Ovary | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Male Genital System | 209.3 | 113.9 | 503.0 | 15 | 0.0 | 0.0 | 146.3 | 0 |
| Prostate | 209.3 | 113.9 | 503.0 | 15 | 0.0 | 0.0 | 146.3 | 0 |
| Testis | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Penis | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2002

| | American Indian/Alaska Native | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 18.1 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 44.7 | 0 | ^ | ^ | ^ | ^ |
| Respiratory System | 89.1 | 59.5 | 145.5 | 30 | 98.3 | 65.5 | 143.8 | 30 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Larynx | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 81.0 | 52.7 | 136.3 | 27 | 96.0 | 63.5 | 141.3 | 29 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Bones and Joints | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 18.1 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 44.7 | 0 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 18.1 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 44.7 | 0 | ^ | ^ | ^ | ^ |
| Breast | 0.0 | 0.0 | 44.7 | 0 | 129.8 | 94.1 | 177.3 | 47 |
| Female Genital System | 0.0 | 0.0 | 44.7 | 0 | 24.2 | 10.5 | 51.3 | 9 |
| Cervix Uteri | 0.0 | 0.0 | 44.7 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 44.7 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 44.7 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Ovary | 0.0 | 0.0 | 44.7 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 44.7 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 44.7 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 100.4 | 62.6 | 166.0 | 30 | 0.0 | 0.0 | 18.1 | 0 |
| Prostate | 90.0 | 53.6 | 155.1 | 25 | 0.0 | 0.0 | 18.1 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 18.1 | 0 |
| Penis | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Other Male Genital Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 18.1 | 0 |
| Urinary System | 38.6 | 20.4 | 87.1 | 14 | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 35.9 | 18.5 | 84.1 | 13 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 44.7 | 0 | ^ | ^ | ^ | ^ |
| Other Urinary Organs | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2002

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 74.3 | 0 |
| Pancreas | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Respiratory System | ^ | ^ | ^ | ^ | 64.4 | 23.1 | 161.6 | 8 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Larynx | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | 64.4 | 23.1 | 161.6 | 8 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Bones and Joints | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Melanoma of the Skin | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Breast | 0.0 | 0.0 | 127.1 | 0 | 88.3 | 48.3 | 177.0 | 15 |
| Female Genital System | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Ovary | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Vulva | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Male Genital System | 215.6 | 99.2 | 427.6 | 12 | 0.0 | 0.0 | 74.3 | 0 |
| Prostate | 209.8 | 94.4 | 421.8 | 11 | 0.0 | 0.0 | 74.3 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 74.3 | 0 |
| Penis | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2002

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Brain and Other Nervous System | 7.7 | 4.6 | 15.9 | 19 | 5.7 | 2.8 | 11.2 | 11 |
| Brain | 7.3 | 4.3 | 15.5 | 18 | 5.7 | 2.8 | 11.2 | 11 |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Endocrine System | 4.6 | 2.1 | 12.7 | 10 | 11.1 | 7.3 | 16.9 | 28 |
| Thyroid | 3.6 | 1.6 | 11.3 | 9 | 10.7 | 7.0 | 16.5 | 27 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphoma | 39.3 | 28.0 | 55.4 | 60 | 20.8 | 14.4 | 29.5 | 39 |
| Hodgkin Lymphoma | 2.2 | 0.8 | 9.7 | 6 | 3.6 | 1.6 | 8.0 | 9 |
| Hodgkin - Nodal | 2.2 | 0.8 | 9.7 | 6 | 3.6 | 1.6 | 8.0 | 9 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 7.7 | 0 | 0.0 | 0.0 | 3.5 | 0 |
| Non-Hodgkin Lymphoma | 37.2 | 26.0 | 53.1 | 54 | 17.2 | 11.3 | 25.5 | 30 |
| NHL - Nodal | 27.3 | 17.9 | 41.5 | 40 | 12.3 | 7.3 | 19.8 | 21 |
| NHL - Extranodal | 9.9 | 4.6 | 20.6 | 14 | 4.9 | 2.2 | 10.2 | 9 |
| Myeloma | 4.2 | 1.7 | 12.3 | 8 | ^ | ^ | ^ | ^ |
| Leukemia | 12.9 | 7.5 | 23.2 | 25 | 9.8 | 5.2 | 17.0 | 14 |
| Lymphocytic Leukemia | 6.4 | 2.4 | 16.1 | 10 | 7.2 | 3.4 | 13.7 | 10 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 5.3 | 1.6 | 14.9 | 7 | 6.2 | 2.7 | 12.7 | 8 |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Myeloid and Monocytic Leukemia | 6.1 | 3.1 | 14.4 | 14 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 3.5 | 1.5 | 11.1 | 9 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 7.7 | 0 | ^ | ^ | ^ | ^ |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 7.7 | 0 | 0.0 | 0.0 | 3.5 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 7.7 | 0 | 0.0 | 0.0 | 3.5 | 0 |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Mesothelioma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.5 | 0 |
| Miscellaneous | 11.1 | 5.6 | 21.9 | 17 | 5.9 | 2.6 | 11.8 | 9 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2002

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Brain | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Endocrine System | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 146.3 | 0 |
| NHL - Extranodal | 0.0 | 0.0 | 286.0 | 0 | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 146.3 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 146.3 | 0 |
| Lymphocytic Leukemia | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 146.3 | 0 |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 146.3 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Other Leukemia | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Mesothelioma | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 286.0 | 0 | 0.0 | 0.0 | 146.3 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2002

| | American Indian/Alaska Native | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 18.1 | 0 |
| Brain | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 18.1 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Endocrine System | 0.0 | 0.0 | 44.7 | 0 | 16.4 | 5.8 | 40.7 | 7 |
| Thyroid | 0.0 | 0.0 | 44.7 | 0 | 16.4 | 5.8 | 40.7 | 7 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Myeloma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Leukemia | 9.5 | 3.2 | 53.0 | 6 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 18.1 | 0 |
| Acute Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 18.1 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 44.7 | 0 | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Other Leukemia | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Mesothelioma | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 44.7 | 0 | 0.0 | 0.0 | 18.1 | 0 |
| Miscellaneous | 27.9 | 10.0 | 79.0 | 7 | 19.0 | 6.9 | 45.0 | 7 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2002

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Brain | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Endocrine System | 0.0 | 0.0 | 127.1 | 0 | 35.0 | 11.9 | 114.5 | 6 |
| Thyroid | 0.0 | 0.0 | 127.1 | 0 | 35.0 | 11.9 | 114.5 | 6 |
| Other Endocrine including Thymus | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 74.3 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 74.3 | 0 |
| NHL - Nodal | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| NHL - Extranodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 74.3 | 0 |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 74.3 | 0 |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 74.3 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 127.1 | 0 | ^ | ^ | ^ | ^ |
| Myeloid and Monocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 74.3 | 0 |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 74.3 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Other Leukemia | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Mesothelioma | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 127.1 | 0 | 0.0 | 0.0 | 74.3 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 74.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2003

| | White | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 491.6 | 454.1 | 532.6 | 852 | 366.2 | 337.3 | 397.3 | 677 |
| Oral Cavity and Pharynx | 14.3 | 8.1 | 24.9 | 25 | 4.1 | 1.4 | 9.5 | 6 |
| Lip | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.3 | 0 |
| Tongue | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.3 | 0 |
| Floor of Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.3 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.3 | 0 |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.3 | 0 |
| Oropharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.3 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.3 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 6.9 | 0 | 0.0 | 0.0 | 3.3 | 0 |
| Digestive System | 92.2 | 75.7 | 112.7 | 154 | 48.7 | 38.2 | 61.5 | 83 |
| Esophagus | 7.0 | 3.8 | 14.8 | 16 | ^ | ^ | ^ | ^ |
| Stomach | 5.3 | 2.3 | 13.2 | 10 | 3.6 | 1.2 | 8.7 | 6 |
| Small Intestine | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon and Rectum | 49.8 | 37.7 | 66.1 | 77 | 26.7 | 19.2 | 36.5 | 47 |
| Colon excluding Rectum | 36.8 | 26.0 | 51.9 | 51 | 15.8 | 10.1 | 23.9 | 27 |
| Cecum | ^ | ^ | ^ | ^ | 4.9 | 1.9 | 10.6 | 7 |
| Appendix | 0.0 | 0.0 | 6.9 | 0 | 0.0 | 0.0 | 3.3 | 0 |
| Ascending Colon | 7.9 | 3.4 | 17.4 | 11 | ^ | ^ | ^ | ^ |
| Hepatic Flexure | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 6.9 | 0 | 0.0 | 0.0 | 3.3 | 0 |
| Descending Colon | 7.7 | 3.5 | 16.8 | 10 | 0.0 | 0.0 | 3.3 | 0 |
| Sigmoid Colon | 10.2 | 5.0 | 20.2 | 14 | 4.3 | 2.0 | 9.1 | 10 |
| Large Intestine, NOS | 5.4 | 1.4 | 15.1 | 7 | ^ | ^ | ^ | ^ |
| Rectum and Rectosigmoid Junction | 13.0 | 8.1 | 22.3 | 26 | 10.9 | 6.4 | 17.8 | 20 |
| Rectosigmoid Junction | 5.3 | 2.4 | 13.1 | 10 | ^ | ^ | ^ | ^ |
| Rectum | 7.7 | 4.0 | 16.0 | 16 | 8.2 | 4.4 | 14.4 | 15 |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | 3.1 | 1.1 | 7.7 | 6 |
| Liver and Intrahepatic Bile Duct | 9.8 | 5.1 | 19.2 | 19 | ^ | ^ | ^ | ^ |
| Liver | 9.8 | 5.1 | 19.2 | 19 | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 6.9 | 0 | ^ | ^ | ^ | ^ |
| Gallbladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2003

| | Black | | | | | | | |
|----------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 443.4 | 258.4 | 787.9 | 24 | 363.4 | 221.8 | 587.3 | 25 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 124.2 | 0 |
| Lip | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Tongue | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Salivary Gland | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 124.2 | 0 |
| Nasopharynx | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Tonsil | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Oropharynx | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Hypopharynx | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Digestive System | 182.4 | 67.9 | 480.2 | 8 | ^ | ^ | ^ | ^ |
| Esophagus | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Stomach | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Colon and Rectum | 138.8 | 42.9 | 424.7 | 6 | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 124.2 | 0 |
| Appendix | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Ascending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 124.2 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Transverse Colon | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Descending Colon | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Sigmoid Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 124.2 | 0 |
| Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 124.2 | 0 |
| Liver | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 124.2 | 0 |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Gallbladder | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2003

| | American Indian/Alaska Native | | | | | | | |
|----------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 556.3 | 460.1 | 675.1 | 150 | 456.2 | 386.2 | 536.9 | 164 |
| Oral Cavity and Pharynx | 31.1 | 15.2 | 75.5 | 11 | ^ | ^ | ^ | ^ |
| Lip | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Tongue | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Salivary Gland | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 40.9 | 0 | ^ | ^ | ^ | ^ |
| Gum and Other Mouth | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Nasopharynx | 18.1 | 6.5 | 60.6 | 6 | ^ | ^ | ^ | ^ |
| Tonsil | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Oropharynx | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Hypopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Other Oral Cavity and Pharynx | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Digestive System | 188.8 | 134.3 | 267.6 | 50 | 123.9 | 88.6 | 170.7 | 43 |
| Esophagus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Stomach | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Colon and Rectum | 106.7 | 65.8 | 173.8 | 28 | 72.2 | 44.9 | 111.7 | 23 |
| Colon excluding Rectum | 84.7 | 46.8 | 150.2 | 19 | 64.8 | 39.3 | 102.6 | 21 |
| Cecum | ^ | ^ | ^ | ^ | 20.0 | 6.9 | 46.8 | 6 |
| Appendix | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hepatic Flexure | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Transverse Colon | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Splenic Flexure | 0.0 | 0.0 | 40.9 | 0 | ^ | ^ | ^ | ^ |
| Descending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Sigmoid Colon | ^ | ^ | ^ | ^ | 27.7 | 13.0 | 54.8 | 10 |
| Large Intestine, NOS | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Rectum and Rectosigmoid Junction | 22.1 | 10.1 | 63.9 | 9 | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectum | 14.5 | 5.3 | 55.5 | 6 | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Gallbladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2003

| | Asian or Pacific Islander | | | | | | | |
|----------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| All Sites | 422.5 | 287.1 | 627.7 | 39 | 250.2 | 175.8 | 363.9 | 40 |
| Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 67.3 | 0 |
| Lip | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Tongue | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Salivary Gland | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Floor of Mouth | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Gum and Other Mouth | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Nasopharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 67.3 | 0 |
| Tonsil | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Oropharynx | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Hypopharynx | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Other Oral Cavity and Pharynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 67.3 | 0 |
| Digestive System | 195.4 | 101.8 | 367.7 | 16 | 47.1 | 18.2 | 125.3 | 7 |
| Esophagus | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Stomach | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Small Intestine | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Colon and Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Colon excluding Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Cecum | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Appendix | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Ascending Colon | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 67.3 | 0 |
| Hepatic Flexure | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Transverse Colon | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Splenic Flexure | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Descending Colon | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Sigmoid Colon | 0.0 | 0.0 | 114.4 | 0 | ^ | ^ | ^ | ^ |
| Large Intestine, NOS | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Rectum and Rectosigmoid Junction | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Rectosigmoid Junction | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 67.3 | 0 |
| Rectum | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Anus, Anal Canal and Anorectum | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Liver and Intrahepatic Bile Duct | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Liver | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Intrahepatic Bile Duct | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Gallbladder | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2003

| | White | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | 12.8 | 6.6 | 23.9 | 19 | 6.4 | 2.9 | 12.4 | 10 |
| Retroperitoneum | 0.0 | 0.0 | 6.9 | 0 | 0.0 | 0.0 | 3.3 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 6.9 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 94.2 | 77.6 | 114.7 | 143 | 58.8 | 46.8 | 73.3 | 90 |
| Nose, Nasal Cavity and Middle Ear | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.3 | 0 |
| Larynx | 5.8 | 2.6 | 13.8 | 10 | ^ | ^ | ^ | ^ |
| Lung and Bronchus | 88.1 | 72.0 | 108.2 | 132 | 57.7 | 45.7 | 72.0 | 87 |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 6.9 | 0 | ^ | ^ | ^ | ^ |
| Bones and Joints | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Soft Tissue including Heart | 4.7 | 1.9 | 12.6 | 9 | ^ | ^ | ^ | ^ |
| Skin excluding Basal and Squamous | 17.8 | 12.5 | 27.0 | 45 | 10.6 | 6.4 | 17.0 | 22 |
| Melanoma of the Skin | 15.8 | 11.1 | 24.4 | 42 | 9.3 | 5.5 | 15.4 | 20 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Breast | ^ | ^ | ^ | ^ | 128.5 | 112.2 | 146.9 | 257 |
| Female Genital System | 0.0 | 0.0 | 6.9 | 0 | 40.9 | 32.0 | 51.9 | 83 |
| Cervix Uteri | 0.0 | 0.0 | 6.9 | 0 | 3.7 | 1.5 | 8.3 | 8 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 6.9 | 0 | 19.9 | 14.0 | 28.1 | 41 |
| Corpus Uteri | 0.0 | 0.0 | 6.9 | 0 | 19.6 | 13.7 | 27.7 | 40 |
| Uterus, NOS | 0.0 | 0.0 | 6.9 | 0 | ^ | ^ | ^ | ^ |
| Ovary | 0.0 | 0.0 | 6.9 | 0 | 14.7 | 9.6 | 22.0 | 30 |
| Vagina | 0.0 | 0.0 | 6.9 | 0 | ^ | ^ | ^ | ^ |
| Vulva | 0.0 | 0.0 | 6.9 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 6.9 | 0 | 0.0 | 0.0 | 3.3 | 0 |
| Male Genital System | 142.7 | 123.5 | 165.6 | 250 | 0.0 | 0.0 | 3.3 | 0 |
| Prostate | 136.1 | 117.2 | 158.7 | 233 | 0.0 | 0.0 | 3.3 | 0 |
| Testis | 5.4 | 2.9 | 12.5 | 14 | 0.0 | 0.0 | 3.3 | 0 |
| Penis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.3 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 6.9 | 0 | 0.0 | 0.0 | 3.3 | 0 |
| Urinary System | 55.5 | 43.1 | 71.9 | 89 | 19.8 | 13.3 | 28.7 | 33 |
| Urinary Bladder | 41.6 | 30.7 | 56.6 | 60 | 10.8 | 6.1 | 17.9 | 17 |
| Kidney and Renal Pelvis | 13.9 | 8.5 | 23.6 | 29 | 8.4 | 4.5 | 14.7 | 15 |
| Ureter | 0.0 | 0.0 | 6.9 | 0 | ^ | ^ | ^ | ^ |
| Other Urinary Organs | 0.0 | 0.0 | 6.9 | 0 | 0.0 | 0.0 | 3.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2003

| | Black | | | | | | | |
|---|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 124.2 | 0 |
| Retroperitoneum | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Other Digestive Organs | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Larynx | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Bones and Joints | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Soft Tissue including Heart | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 124.2 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Breast | ^ | ^ | ^ | ^ | 107.8 | 42.3 | 264.2 | 9 |
| Female Genital System | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Cervix Uteri | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Ovary | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Vulva | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Male Genital System | 161.8 | 58.9 | 448.7 | 8 | 0.0 | 0.0 | 124.2 | 0 |
| Prostate | 161.8 | 58.9 | 448.7 | 8 | 0.0 | 0.0 | 124.2 | 0 |
| Testis | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Penis | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 124.2 | 0 |
| Kidney and Renal Pelvis | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2003

| | American Indian/Alaska Native | | | | | | | |
|---|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 40.9 | 0 | ^ | ^ | ^ | ^ |
| Other Digestive Organs | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Respiratory System | 137.6 | 91.8 | 208.5 | 35 | 61.0 | 36.6 | 97.6 | 20 |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Lung and Bronchus | 130.1 | 85.2 | 200.3 | 32 | 61.0 | 36.6 | 97.6 | 20 |
| Trachea, Mediastinum and Other Respiratory Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Bones and Joints | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Skin excluding Basal and Squamous | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Melanoma of the Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Other Non-Epithelial Skin | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Breast | 0.0 | 0.0 | 40.9 | 0 | 153.3 | 116.3 | 201.0 | 61 |
| Female Genital System | 0.0 | 0.0 | 40.9 | 0 | 33.7 | 16.0 | 64.4 | 11 |
| Cervix Uteri | 0.0 | 0.0 | 40.9 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 40.9 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 40.9 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 40.9 | 0 | ^ | ^ | ^ | ^ |
| Ovary | 0.0 | 0.0 | 40.9 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Vulva | 0.0 | 0.0 | 40.9 | 0 | ^ | ^ | ^ | ^ |
| Other Female Genital Organs | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Male Genital System | 90.2 | 53.8 | 152.9 | 25 | 0.0 | 0.0 | 16.7 | 0 |
| Prostate | 80.1 | 45.1 | 142.3 | 20 | 0.0 | 0.0 | 16.7 | 0 |
| Testis | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Penis | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Urinary System | 69.3 | 36.0 | 130.7 | 16 | 18.1 | 7.0 | 41.8 | 7 |
| Urinary Bladder | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Kidney and Renal Pelvis | 36.1 | 15.7 | 85.3 | 10 | 18.1 | 7.0 | 41.8 | 7 |
| Ureter | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2003

| | Asian or Pacific Islander | | | | | | | |
|---|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Other Biliary | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Pancreas | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Retroperitoneum | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Peritoneum, Omentum and Mesentery | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Other Digestive Organs | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 67.3 | 0 |
| Respiratory System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Nose, Nasal Cavity and Middle Ear | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Larynx | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 67.3 | 0 |
| Lung and Bronchus | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Trachea, Mediastinum and Other Respiratory Organs | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Bones and Joints | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Soft Tissue including Heart | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Skin excluding Basal and Squamous | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Melanoma of the Skin | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Other Non-Epithelial Skin | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Breast | 0.0 | 0.0 | 114.4 | 0 | 66.2 | 33.2 | 145.0 | 12 |
| Female Genital System | 0.0 | 0.0 | 114.4 | 0 | 45.7 | 18.9 | 121.1 | 8 |
| Cervix Uteri | 0.0 | 0.0 | 114.4 | 0 | ^ | ^ | ^ | ^ |
| Corpus and Uterus, NOS | 0.0 | 0.0 | 114.4 | 0 | ^ | ^ | ^ | ^ |
| Corpus Uteri | 0.0 | 0.0 | 114.4 | 0 | ^ | ^ | ^ | ^ |
| Uterus, NOS | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Ovary | 0.0 | 0.0 | 114.4 | 0 | ^ | ^ | ^ | ^ |
| Vagina | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Vulva | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Other Female Genital Organs | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Male Genital System | 119.9 | 56.7 | 262.1 | 11 | 0.0 | 0.0 | 67.3 | 0 |
| Prostate | 119.9 | 56.7 | 262.1 | 11 | 0.0 | 0.0 | 67.3 | 0 |
| Testis | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Penis | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Other Male Genital Organs | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Urinary System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Urinary Bladder | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Kidney and Renal Pelvis | 0.0 | 0.0 | 114.4 | 0 | ^ | ^ | ^ | ^ |
| Ureter | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Other Urinary Organs | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2003

| | White | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Brain and Other Nervous System | 7.4 | 4.0 | 15.4 | 17 | 6.1 | 2.9 | 11.9 | 11 |
| Brain | 7.0 | 3.7 | 14.9 | 16 | 5.7 | 2.6 | 11.4 | 10 |
| Cranial Nerves Other Nervous System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Endocrine System | 4.7 | 2.4 | 11.9 | 12 | 12.7 | 8.5 | 18.9 | 30 |
| Thyroid | 3.7 | 1.7 | 10.7 | 10 | 12.7 | 8.5 | 18.9 | 30 |
| Other Endocrine including Thymus | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.3 | 0 |
| Lymphoma | 26.4 | 18.6 | 38.2 | 51 | 14.9 | 9.5 | 22.7 | 27 |
| Hodgkin Lymphoma | 5.5 | 2.7 | 13.2 | 13 | ^ | ^ | ^ | ^ |
| Hodgkin - Nodal | 5.5 | 2.7 | 13.2 | 13 | ^ | ^ | ^ | ^ |
| Hodgkin - Extranodal | 0.0 | 0.0 | 6.9 | 0 | 0.0 | 0.0 | 3.3 | 0 |
| Non-Hodgkin Lymphoma | 20.8 | 13.8 | 32.1 | 38 | 13.1 | 8.0 | 20.6 | 23 |
| NHL - Nodal | 11.6 | 7.0 | 20.5 | 24 | 8.3 | 4.5 | 14.4 | 16 |
| NHL - Extranodal | 9.2 | 4.4 | 19.0 | 14 | 4.8 | 1.8 | 10.6 | 7 |
| Myeloma | 2.4 | 0.8 | 9.4 | 6 | 4.8 | 1.9 | 10.5 | 7 |
| Leukemia | 9.5 | 5.7 | 17.6 | 21 | 5.0 | 2.0 | 10.6 | 8 |
| Lymphocytic Leukemia | 4.1 | 1.9 | 11.3 | 10 | ^ | ^ | ^ | ^ |
| Acute Lymphocytic Leukemia | 2.2 | 0.8 | 9.1 | 6 | ^ | ^ | ^ | ^ |
| Chronic Lymphocytic Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.3 | 0 |
| Myeloid and Monocytic Leukemia | 4.5 | 1.9 | 12.0 | 9 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 2.6 | 1.0 | 9.6 | 6 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 6.9 | 0 | 0.0 | 0.0 | 3.3 | 0 |
| Chronic Myeloid Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 6.9 | 0 | 0.0 | 0.0 | 3.3 | 0 |
| Other Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Acute Leukemia | 0.0 | 0.0 | 6.9 | 0 | 0.0 | 0.0 | 3.3 | 0 |
| Aleukemic, Subleukemic and NOS | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Mesothelioma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 3.3 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 6.9 | 0 | 0.0 | 0.0 | 3.3 | 0 |
| Miscellaneous | 13.4 | 7.5 | 23.9 | 21 | 8.5 | 4.4 | 15.0 | 14 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2003

| | Black | | | | | | | |
|-------------------------------------|-------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Brain | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Endocrine System | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Lymphoma | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Hodgkin Lymphoma | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Non-Hodgkin Lymphoma | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| NHL - Nodal | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| NHL - Extranodal | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Myeloma | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Leukemia | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 248.7 | 0 | ^ | ^ | ^ | ^ |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Other Leukemia | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Mesothelioma | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |
| Miscellaneous | 0.0 | 0.0 | 248.7 | 0 | 0.0 | 0.0 | 124.2 | 0 |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2003

| | American Indian/Alaska Native | | | | | | | |
|-------------------------------------|-------------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Brain and Other Nervous System | 0.0 | 0.0 | 40.9 | 0 | ^ | ^ | ^ | ^ |
| Brain | 0.0 | 0.0 | 40.9 | 0 | ^ | ^ | ^ | ^ |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Endocrine System | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Thyroid | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Hodgkin - Nodal | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Non-Hodgkin Lymphoma | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Nodal | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| NHL - Extranodal | 0.0 | 0.0 | 40.9 | 0 | ^ | ^ | ^ | ^ |
| Myeloma | 0.0 | 0.0 | 40.9 | 0 | ^ | ^ | ^ | ^ |
| Leukemia | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Acute Lymphocytic Leukemia | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 16.7 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 40.9 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 40.9 | 0 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Other Leukemia | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Mesothelioma | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Kaposi Sarcoma | 0.0 | 0.0 | 40.9 | 0 | 0.0 | 0.0 | 16.7 | 0 |
| Miscellaneous | 19.8 | 7.1 | 63.2 | 6 | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.

Malignant Cancer Cases in Alaska, Diagnosis Years 1996 to 2003, by Race and Sex

2003

| | Asian or Pacific Islander | | | | | | | |
|-------------------------------------|---------------------------|----------|----------|-------|--------|----------|----------|-------|
| | Male | | | | Female | | | |
| | Rate | Lower CI | Upper CI | Count | Rate | Lower CI | Upper CI | Count |
| Eye and Orbit | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Brain and Other Nervous System | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 67.3 | 0 |
| Brain | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 67.3 | 0 |
| Cranial Nerves Other Nervous System | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Endocrine System | 0.0 | 0.0 | 114.4 | 0 | ^ | ^ | ^ | ^ |
| Thyroid | 0.0 | 0.0 | 114.4 | 0 | ^ | ^ | ^ | ^ |
| Other Endocrine including Thymus | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Lymphoma | 0.0 | 0.0 | 114.4 | 0 | ^ | ^ | ^ | ^ |
| Hodgkin Lymphoma | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Hodgkin - Nodal | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Hodgkin - Extranodal | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Non-Hodgkin Lymphoma | 0.0 | 0.0 | 114.4 | 0 | ^ | ^ | ^ | ^ |
| NHL - Nodal | 0.0 | 0.0 | 114.4 | 0 | ^ | ^ | ^ | ^ |
| NHL - Extranodal | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Myeloma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 67.3 | 0 |
| Leukemia | 0.0 | 0.0 | 114.4 | 0 | ^ | ^ | ^ | ^ |
| Lymphocytic Leukemia | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Acute Lymphocytic Leukemia | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Chronic Lymphocytic Leukemia | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Other Lymphocytic Leukemia | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Myeloid and Monocytic Leukemia | 0.0 | 0.0 | 114.4 | 0 | ^ | ^ | ^ | ^ |
| Acute Myeloid Leukemia | 0.0 | 0.0 | 114.4 | 0 | ^ | ^ | ^ | ^ |
| Acute Monocytic Leukemia | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Chronic Myeloid Leukemia | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Other Myeloid/Monocytic Leukemia | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Other Leukemia | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Other Acute Leukemia | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Aleukemic, Subleukemic and NOS | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Mesothelioma | 0.0 | 0.0 | 114.4 | 0 | 0.0 | 0.0 | 67.3 | 0 |
| Kaposi Sarcoma | ^ | ^ | ^ | ^ | 0.0 | 0.0 | 67.3 | 0 |
| Miscellaneous | ^ | ^ | ^ | ^ | ^ | ^ | ^ | ^ |

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard; Confidence intervals are 95% for rates.

~ Statistic could not be calculated.

^ Statistic not displayed due to fewer than 6 cases.